

# phostensin (C-1): sc-365398

## BACKGROUND

In eukaryotes, the phosphorylation and dephosphorylation of proteins on serine and threonine residues is an essential means of regulating a broad range of cellular functions, including division, homeostasis and apoptosis. A group of proteins that are intimately involved in this process are the protein phosphatases, specifically PP1 (protein phosphatase 1), which is targeted to different substrates throughout the cell. Phostensin, also known as KIAA1949, is a 613 amino acid protein that localizes to both the cytoplasm and the cytoskeleton. Expressed predominately in spleen, ovary, lung and liver tissue, phostensin functions as a regulatory subunit that interacts with and targets PP1 to F-Actin in the cytoskeleton. Two isoforms of phostensin exist due to alternative splicing events.

## REFERENCES

1. Nagase, T., Kikuno, R. and Ohara, O. 2001. Prediction of the coding sequences of unidentified human genes. XXII. The complete sequences of 50 new cDNA clones which code for large proteins. *DNA Res.* 8: 319-327.
2. Terry-Lorenzo, R.T., Elliott, E., Weiser, D.C., Prickett, T.D., Brautigan, D.L. and Shenolikar, S. 2002. Neurabins recruit protein phosphatase-1 and inhibitor-2 to the Actin cytoskeleton. *J. Biol. Chem.* 277: 46535-46543.
3. Oliver, C.J., Terry-Lorenzo, R.T., Elliott, E., Bloomer, W.A., Li, S., Brautigan, D.L., Colbran, R.J. and Shenolikar, S. 2002. Targeting protein phosphatase 1 (PP1) to the Actin cytoskeleton: the neurabin I/PP1 complex regulates cell morphology. *Mol. Cell. Biol.* 22: 4690-4701.
4. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 610990. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Olsen, J.V., Blagoev, B., Gnad, F., Macek, B., Kumar, C., Mortensen, P. and Mann, M. 2006. Global, *in vivo*, and site-specific phosphorylation dynamics in signaling networks. *Cell* 127: 635-648.
6. Kao, S.C., Chen, C.Y., Wang, S.L., Yang, J.J., Hung, W.C., Chen, Y.C., Lai, N.S., Liu, H.T., Huang, H.L., Chen, H.C., Lin, T.H. and Huang, H.B. 2007. Identification of phostensin, a PP1 F-Actin cytoskeleton targeting subunit. *Biochem. Biophys. Res. Commun.* 356: 594-598.

## CHROMOSOMAL LOCATION

Genetic locus: PPP1R18 (human) mapping to 6p21.33.

## SOURCE

phostensin (C-1) is a mouse monoclonal antibody raised against amino acids 121-420 mapping within an internal region of phostensin of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>2b</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

phostensin (C-1) is recommended for detection of phostensin of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for phostensin siRNA (h): sc-95055, phostensin shRNA Plasmid (h): sc-95055-SH and phostensin shRNA (h) Lentiviral Particles: sc-95055-V.


Molecular Weight of phostensin isoforms: 26/68 kDa.

Positive Controls: MIA PaCa-2 whole cell lysate: sc-2285 or Hep G2 cell lysate: sc-2227.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

## DATA



phostensin (C-1): sc-365398. Western blot analysis of phostensin expression in MIA PaCa-2 (A) and Hep G2 (B) whole cell lysates.



phostensin (C-1): sc-365398. Immunoperoxidase staining of formalin fixed, paraffin-embedded human lymph node tissue showing cytoplasmic staining of cells in germinal and non-germinal centers.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.