

SYP (C-12): sc-365447

BACKGROUND

Synaptic vesicles participate in a cycle of fusion with the plasma membrane and reformation by endocytosis. Synaptic vesicle protein synaptophysin (SYP) is targeted to early endosomes in transfected fibroblasts and in neuroendocrine cells. SYP is an N-glycosylated integral membrane protein found in neurons and endocrine cells that associates into hexamers to form a large conductance channel. SYP contains four transmembrane domains and may function as a gap junction-like channel. Membrane cholesterol specifically interacts with SYP to play a role in vesicle formation. Synaptobrevin (VAMP) also binds to SYP and the resultant complex is upregulated during neuronal development, but is absent in exocytosis fusion complex. Thus, the synaptophysin-synaptobrevin complex is not essential for exocytosis, but rather provides a pool of synaptobrevin for exocytosis. In addition, the tail domain of brain Myosin V also forms a stable complex with synaptobrevin II and SYP, and this complex is disassembled upon the depolarization-induced entry of Ca²⁺ into intact nerve endings.

CHROMOSOMAL LOCATION

Genetic locus: SYP (human) mapping to Xp11.23; Syp (mouse) mapping to X A1.1.

SOURCE

SYP (C-12) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 289-312 at the C-terminus of SYP of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-365447 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

SYP (C-12) is recommended for detection of SYP of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for SYP siRNA (h): sc-36597, SYP siRNA (m): sc-36596, SYP shRNA Plasmid (h): sc-36597-SH, SYP shRNA Plasmid (m): sc-36596-SH, SYP shRNA (h) Lentiviral Particles: sc-36597-V and SYP shRNA (m) Lentiviral Particles: sc-36596-V.

Molecular Weight of SYP: 38-48 kDa.

Positive Controls: IMR-32 cell lysate: sc-2409, MDA-MB-231 cell lysate: sc-2232 or mouse brain extract: sc-2253.


RESEARCH USE

For research use only, not for use in diagnostic procedures.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



SYP (C-12): sc-365447. Western blot analysis of SYP expression in IMR-32 (A) and MDA-MB-231 (B) whole cell lysates.



SYP (C-12): sc-365447. Immunoperoxidase staining of formalin fixed, paraffin-embedded human cerebral cortex (A) and mouse brain (B) tissue showing neuropil staining.

SELECT PRODUCT CITATIONS


1. Khan, M., Rutten, B.P.F. and Kim, M.O. 2019. MST1 regulates neuronal cell death via JNK/Casp3 signaling pathway in HFD mouse brain and HT22 cells. *Int. J. Mol. Sci.* 20: 2504.
2. Bektas, H., Algul, S., Altindag, F., Yegin, K., Akdag, Z. and Dasdag, S. 2022. Effects of 3.5 GHz (5G) radiofrequency radiation on ghrelin, nesfatin-1, and irisin levels in diabetic and healthy brains. *J. Chem. Neuroanat.* E-published.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **SYP (D-4): sc-17750** for SYP antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.