# HCN3 (B-12): sc-365451



The Boures to Overtion

#### **BACKGROUND**

Hyperpolarization-activated, cyclic nucleotide-binding channels (HCN) are voltage-gated cation channels that are activated by direct binding of intracellular cyclic nucleotides. The HCN family consists of four members (HCN1-4), each with a core transmembrane segment domain and a C-terminal 120 amino-acid cyclic nucleotide-binding domain motif. HCN channels are expressed in the brain, heart, thalamus and testis. The pacemaker properties of HCN channels contribute to spontaneous rhythmic activity in the brain and heart. HCN3 contains a segment characterized by a series of positively charged amino acids at every third position. This region designated S4 is likely to be the voltage sensor of the protein. In the brain, HCN3 and HCN4 exhibit subcortical distribution mainly concentrated in the hypothalamus and thalamus, respectively.

### **REFERENCES**

- 1. Notomi, T., et al. 2004. Immunohistochemical localization of Ih channel subunits, HCN1-4, in the rat brain. J. Comp. Neurol. 471: 241-276.
- Bajorat, R., et al. 2005. Functional significance of HCN2/3-mediated I<sub>h</sub> in striatal cells at early developmental stages. J. Neurosci. Res. 82: 206-213.
- 3. Stieber, J., et al. 2005. Functional expression of the human HCN3 channel. J. Biol. Chem. 280: 34635-34643.
- Varghese, A., et al. 2005. Endogenous channels in HEK cells and potential roles in HCN ionic current measurements. Prog. Biophys. Mol. Biol. 90: 26-37.
- Yamada, R., et al. 2005. Hyperpolarization-activated cyclic nucleotide-gated cation channels regulate auditory coincidence detection in nucleus laminaris of the chick. J. Neurosci. 25: 8867-8877.
- SWISS-PROT/TrEMBL (Q9P1Z3). World Wide Web URL: http://www.expasy.ch/sprot/sprot-top.html

## CHROMOSOMAL LOCATION

Genetic locus: HCN3 (human) mapping to 1q22.

#### **SOURCE**

HCN3 (B-12) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 757-778 within a C-terminal cytoplasmic domain of HCN3 of human origin.

# **PRODUCT**

Each vial contains 200  $\mu$ g lgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-365451 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

#### **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

#### **APPLICATIONS**

HCN3 (B-12) is recommended for detection of HCN3 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for HCN3 siRNA (h): sc-45651, HCN3 shRNA Plasmid (h): sc-45651-SH and HCN3 shRNA (h) Lentiviral Particles: sc-45651-V.


Molecular Weight of HCN3: 86 kDa.

Positive Controls: HCN3 (h): 293T Lysate: sc-114783.

# **RECOMMENDED SUPPORT REAGENTS**

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-lgG $\kappa$  BP-HRP: sc-516102 or m-lgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>TM</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-lgG $\kappa$  BP-FITC: sc-516140 or m-lgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

#### **DATA**



HCN3 (B-12): sc-365451. Western blot analysis of HCN3 expression in non-transfected: sc-117752 (A) and human HCN3 transfected: sc-114783 (B) 293T whole cell bycates

## **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

#### **PROTOCOLS**

See our web site at www.scbt.com for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com