

p-PDGFR- β (H-3): sc-365465

BACKGROUND

Platelet derived growth factor (PDGF) is a mitogen for mesenchyme- and glia-derived cells. PDGF consists of two chains, A and B, which dimerize to form functionally distinct isoforms, PDGF-AA, PDGF-AB, and PDGF-BB. These three isoforms bind with different affinities to two receptor types, α and β , which are endowed with protein tyrosine kinase domains and undergo either homo- or heterodimerization as a consequence of ligand binding. Ligand stimulation of PDGFR- β leads to autophosphorylation at Tyr-857, which is the major autophosphorylation site, and Tyr-751, which is the major *in vitro* phosphorylation site. Autophosphorylation of Tyr-751, which lies in the kinase insert region, is required for binding of phosphatidylinositol-3 kinase to the receptor. These autophosphorylation events largely contribute to signal transduction through the PDGF receptor.

REFERENCES

- Ross, R., et al. 1986. The biology of platelet-derived growth factor. *Cell* 46: 155-169.
- Hart, C.E., et al. 1988. Two classes of PDGF receptor recognize different isoforms of PDGF. *Science* 240: 1529-1531.
- Heldin, C., et al. 1988. Binding of different dimeric forms of PDGF to human fibroblasts: evidence for two separate receptor types. *EMBO J.* 7: 1387-1393.
- Kazlauskas, A., et al. 1989. Autophosphorylation of the PDGF receptor in the kinase insert region regulates interactions with cell proteins. *Cell* 58: 1121-1133.
- Kelly, J.D., et al. 1991. Platelet-derived growth factor (PDGF) stimulates PDGF receptor subunit dimerization and intersubunit *trans*-phosphorylation. *J. Biol. Chem.* 266: 8987-8992.
- Nishimura, R., et al. 1993. Two signaling molecules share a phosphotyrosine-containing binding site in the platelet-derived growth factor receptor. *Mol. Cell. Biol.* 13: 6889-6896.

CHROMOSOMAL LOCATION

Genetic locus: PDGFRB (human) mapping to 5q32; Pdgfrb (mouse) mapping to 18 E1.

SOURCE

p-PDGFR- β (H-3) is a mouse monoclonal antibody specific for an epitope corresponding to a short amino acid sequence containing Tyr 716 phosphorylated PDGFR- β of human origin.

PRODUCT

Each vial contains 200 μ g IgG₃ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-365465 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

p-PDGFR- β (H-3) is recommended for detection of p-PDGFR- β of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for PDGFR- β siRNA (h): sc-29442, PDGFR- β siRNA (m): sc-36200, PDGFR- β shRNA Plasmid (h): sc-29442-SH, PDGFR- β shRNA Plasmid (m): sc-36200-SH, PDGFR- β shRNA (h) Lentiviral Particles: sc-29442-V and PDGFR- β shRNA (m) Lentiviral Particles: sc-36200-V.



Molecular Weight of p-PDGFR- β : 190 kDa.

Positive Controls: NIH/3T3 whole cell lysate: sc-2210 or CCD-1064Sk + PDGF cell lysate: sc-2264.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



Western blot analysis of PDGFR- β phosphorylation in non-transfected (A, D), untreated human PDGFR- β transfected: sc-159386 (B, E) and lambda protein phosphatase (sc-200312A) treated human PDGFR- β transfected: sc-159386 (C, F) 293T whole cell lysates. Antibodies tested include p-PDGFR- β (H-3): sc-365465 (A, B, C) and PDGFR- β (11H4): sc-80991 (D, E, F).

Western blot analysis of PDGFR- β phosphorylation in non-transfected (A, D), untreated human PDGFR- β transfected: sc-159386 (B, E) and lambda protein phosphatase (sc-200312A) treated human PDGFR- β transfected: sc-159386 (C, F) 293T whole cell lysates. Antibodies tested include p-PDGFR- β (H-3): sc-365465 (A, B, C) and PDGFR- β (11H4): sc-80991 (D, E, F).

SELECT PRODUCT CITATIONS

- Panchbhai, N., et al. 2021. P68 RNA helicase facilitates breast cancer progression by promoting proliferation and migration via PDGFR- β /AR axis. *J. Cancer* 12: 6543-6552.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.