SANTA CRUZ BIOTECHNOLOGY, INC.

ZAP-70 (A-1): sc-365490



BACKGROUND

The activation of T lymphocytes by antigens is mediated by the T cell receptor (TCR) which is a multisubunit complex assembled from at least six different genes. The TCR subunits include the Ti α and β chains, the CD3 γ , δ and ε chains and a ζ -containing homodimer or heterodimer. The disulfide-linked Ti α - β heterodimer is responsible for antigen recognition, but the short five amino acid cytoplasmic domains of Ti α and β are unlikely to be sufficient to couple to intracellular signaling pathways. In contrast, the structured features of the CD3 and ζ subunits suggest a role in signal transduction. Of these, the ζ chain, which is expressed as either a homodimer or heterodimer, has a short extracellular domain of only nine amino acids, but a larger 113 amino acid cytoplasmic domain. A tyrosine phosphoprotein, ZAP-70, has been identified that associates with ζ and undergoes tyrosine phosphorylation following TCR stimulation.

CHROMOSOMAL LOCATION

Genetic locus: ZAP70 (human) mapping to 2q11.2; Zap70 (mouse) mapping to 1 B.

SOURCE

ZAP-70 (A-1) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 5-35 near the N-terminus of ZAP-70 of human origin.

PRODUCT

Each vial contains 200 μg lgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-365490 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

ZAP-70 (A-1) is recommended for detection of ZAP-70 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

ZAP-70 (A-1) is also recommended for detection of ZAP-70 in additional species, including bovine.

Suitable for use as control antibody for ZAP-70 siRNA (h): sc-29526, ZAP-70 siRNA (m): sc-36867, ZAP-70 shRNA Plasmid (h): sc-29526-SH, ZAP-70 shRNA Plasmid (m): sc-36867-SH, ZAP-70 shRNA (h) Lentiviral Particles: sc-29526-V and ZAP-70 shRNA (m) Lentiviral Particles: sc-36867-V.

Molecular Weight of ZAP-70: 70 kDa.

Positive Controls: CCRF-CEM cell lysate: sc-2225, BYDP whole cell lysate: sc-364368 or MOLT-4 cell lysate: sc-2233.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA





ZAP-70 (A-1): sc-365490. Western blot analysis of ZAP-70 expression in CCRF-CEM (A), MOLT-4 (B) and BYDP (C) whole cell lysates.

ZAP-70 (A-1): sc-365490. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization

SELECT PRODUCT CITATIONS

 Ivanova, E. and Carpino, N. 2016. Negative regulation of TCR signaling by ubiquitination of ZAP-70 Lys-217. Mol. Immunol. 73: 19-28.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **ZAP-70 (1E7.2): sc-32760** for ZAP-70 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor[®] 488, 546, 594, 647, 680 and 790.