

p-VASP (A-7): sc-365563

BACKGROUND

The Wiskott-Aldrich syndrome (WAS) is characterized by thrombocytopenia, eczema, defects in cell-mediated and humoral immunity and a propensity for lymphoproliferative diseases. The syndrome is the result of a mutation in the gene encoding a proline-rich protein termed WASP. A distantly related protein, VASP (vasodilator-stimulated phosphoprotein), is involved in the maintenance of cytoarchitecture by interacting with Actin-like filaments. VASP shares a limited degree of homology with the amino-terminus of WASP, which is frequently mutated in WAS patients. An established substrate of cAMP and cGMP dependent kinases, VASP is phosphorylated on a regulatory serine residue 157 and localizes to focal adhesions, microfilaments and highly active regions of the plasma membrane. VASP is also phosphorylated on serine 239 by cGMP-dependent protein kinase.

REFERENCES

1. Reinhard, M., et al. 1992. The 46/50 kDa phosphoprotein VASP purified from human platelets is a novel protein associated with Actin filaments and focal contacts. *EMBO J.* 11: 2063-2070.
2. Butt, E., et al. 1994. cAMP- and cGMP-dependent protein kinase phosphorylation sites of the focal adhesion vasodilator-stimulated phosphoprotein (VASP) *in vitro* and in intact human platelets. *J. Biol. Chem.* 269: 14509-14517.
3. Reinhard, M., et al. 1995. Identification, purification and characterization of a Zyxin-related protein that binds the focal adhesion and microfilament protein VASP (vasodilator-stimulated phosphoprotein). *Proc. Natl. Acad. Sci. USA* 92: 7956-7960.
4. Remold-O'Donnell, E., et al. 1996. Defects in Wiskott-Aldrich syndrome blood cells. *Blood* 87: 2621-2631.
5. Stewart, D.M., et al. 1996. Studies of the expression of the Wiskott-Aldrich syndrome protein. *J. Clin. Invest.* 97: 2627-2634.

CHROMOSOMAL LOCATION

Genetic locus: VASP (human) mapping to 19q13.32; Vasp (mouse) mapping to 7 A3.

SOURCE

p-VASP (A-7) is a mouse monoclonal antibody epitope corresponding to a short amino acid sequence containing Ser 157 phosphorylated VASP of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-365563 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

p-VASP (A-7) is recommended for detection of Ser 157 phosphorylated VASP of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for VASP siRNA (h): sc-29516, VASP siRNA (m): sc-36809, VASP shRNA Plasmid (h): sc-29516-SH, VASP shRNA Plasmid (m): sc-36809-SH, VASP shRNA (h) Lentiviral Particles: sc-29516-V and VASP shRNA (m) Lentiviral Particles: sc-36809-V.


Molecular Weight of p-VASP: 50 kDa.

Positive Controls: VASP (h): 293T Lysate: sc-114829, HUV-EC-C whole cell lysate: sc-364180 or MTE1D whole cell lysate: sc-364918.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto B Blocking Reagent: sc-2335 (use 50 mM NaF, sc-24988, as diluent), Lambda Phosphatase: sc-200312A and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



Western blot analysis of VASP phosphorylation in non-transfected: sc-117752 (A,D), untreated human VASP transfected: sc-114829 (B,E) and lambda protein phosphatase (sc-200312A) treated human VASP transfected: sc-114829 (C,F) 293T whole cell lysates. Antibodies tested include p-VASP (A-7): sc-365563 (A,B,C) and VASP (A-11): sc-46668 (D,E,F).



p-VASP (A-7): sc-365563. Western blot analysis of VASP phosphorylation in HUV-EC-C (A) and MTE1D (B) whole cell lysates.

SELECT PRODUCT CITATIONS

1. Satti, H.H., et al. 2021. Antiplatelet activity of astaxanthin in control- and high cholesterol-fed rats mediated by down-regulation of P2Y₁₂, inhibition of NFκB, and increasing intracellular levels of cAMP. *Platelets* 32: 469-478.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.