SANTA CRUZ BIOTECHNOLOGY, INC.

TAP (D-9): sc-365576



BACKGROUND

The constitutive transport element (CTE) of type D retroviruses serves as a signal of nuclear export for unspliced viral RNAs. TAP (also known as NXF1) mediates the export of CTE-containing simian type D retroviral RNAs through binding directly to the CTE. TAP is associated with a recognized mRNA export pathway and is a member of the multigene family of NXF proteins. NXF proteins belong to an evolutionarily conserved family of proteins, which are characterized by a leucine-rich-repeat domain (LRR) followed by a region known as the nuclear transport factor 2 (NTF2)-like domain.

REFERENCES

- 1. Tan, W., et al. 2000. The mRNA export in *Caenorhabditis elegans* is mediated by Ce-NXF-1, an ortholog of human TAP/NXF and *Saccharomyces cerevisiae* Mex67p. RNA 6: 1762-1772.
- 2. Herold, A., et al. 2000. TAP (NXF1) belongs to a multigene family of putative RNA export factors with a conserved modular architecture. Mol. Cell. Biol. 20: 8996-9008.

CHROMOSOMAL LOCATION

Genetic locus: NXF1 (human) mapping to 11q12.3; Nxf1 (mouse) mapping to 19 A.

SOURCE

TAP (D-9) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 57-85 near the N-terminus of TAP of human origin.

PRODUCT

Each vial contains 200 μg lgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-365576 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

TAP (D-9) is recommended for detection of TAP of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for TAP siRNA (h): sc-38142, TAP siRNA (m): sc-38143, TAP shRNA Plasmid (h): sc-38142-SH, TAP shRNA Plasmid (m): sc-38143-SH, TAP shRNA (h) Lentiviral Particles: sc-38142-V and TAP shRNA (m) Lentiviral Particles: sc-38143-V.

Molecular Weight of TAP: 73 kDa.

Positive Controls: Jurkat nuclear extract: sc-2132, HeLa nuclear extract: sc-2120 or K-562 whole cell lysate: sc-2203.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG K BP-HRP: sc-516102 or m-IgG K BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG K BP-FITC: sc-516140 or m-IgG K BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA





TAP (D-9): sc-365576. Western blot analysis of TAP expression in K-562 whole cell lysate (**A**) and MOLT-4 (**B**), Jurkat (**C**) and HeLa (**D**) nuclear extracts. TAP (D-9): sc-365576. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic and nuclear localization.

SELECT PRODUCT CITATIONS

 Ke, H., et al. 2019. Porcine reproductive and respiratory syndrome virus nsp1-β protein interacts with nucleoporin 62 (Nup62) to promote viral replication and immune evasion. J. Virol. 93: e00469-19.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.