

IGF2BP3 (C-11): sc-365641

BACKGROUND

Insulin like growth factor 2 mRNA binding proteins (IGF2BPs) bind RNA and influence RNA synthesis and metabolism. IGF2BP1, also known as coding region determinant-binding protein/insulin-like growth factor II mRNA-binding protein (CRD-BP), IMP1 or VICKZ1; IGF2BP2 (IMP2, VICKZ2, p62); and IGF2BP3 (IMP3, KOC1, VICKZ3) contain a unique combination of RNA recognition motifs and four hnRNP K homology domains. IGF2BP1 is abundant in embryonal tissues and is expressed in 81% of colon cancers, 73% of sarcomas and 58.5% of breast cancers. It recognizes c-Myc, IGF-II and t mRNAs, and H19 RNA, and plays a major role in proliferation of K-562 cells by an IGF-II-dependent mechanism. IGF2BP2 binds the 5' UTR of IGF-II mRNA and influences tumor cell growth, in which IGF2BP2 is associated with apoptosis induced by tretinoin. IGF2BP3 knockdown by RNA interference decreases levels of IGF-II protein without affecting IGF-II, c-Myc, or β Actin mRNA and H19 RNA levels. IGF2BP3 is a marker for carcinomas and high-grade dysplastic lesions of pancreatic ductal epithelium.

CHROMOSOMAL LOCATION

Genetic locus: IGF2BP3 (human) mapping to 7p15.3.

SOURCE

IGF2BP3 (C-11) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 149-180 near the N-terminus of IGF2BP3 of human origin.

PRODUCT

Each vial contains 200 μ g IgG κ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

IGF2BP3 (C-11) is available conjugated to Alexa Fluor® 790 (sc-365641 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-365641 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA-

APPLICATIONS

IGF2BP3 (C-11) is recommended for detection of IGF2BP3 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for IGF2BP3 siRNA (h): sc-60846, IGF2BP3 shRNA Plasmid (h): sc-60846-SH and IGF2BP3 shRNA (h) Lentiviral Particles: sc-60846-V.


Molecular Weight of IGF2BP3: 69 kDa.

Positive Controls: IGF2BP3 (h): 293T Lysate: sc-117068, K-562 whole cell lysate: sc-2203 or NTERA-2 cl.D1 whole cell lysate: sc-364181.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



IGF2BP3 (C-11): sc-365641. Western blot analysis of IGF2BP3 expression in non-transfected: sc-117752 (A) and human IGF2BP3 transfected: sc-117068 (B) 293T whole cell lysates and full length human recombinant IGF2BP3 (C). Detection reagent used: m-IgG κ BP-HRP: sc-525408.



IGF2BP3 (C-11): sc-365641. Immunofluorescence staining of formalin-fixed Hep G2 cells showing cytoplasmic localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human small intestine tissue showing cytoplasmic staining of glandular cells (B).

SELECT PRODUCT CITATIONS

- Hu, Y., et al. 2020. METTL3 regulates the malignancy of cervical cancer via post-transcriptional regulation of RAB2B. *Eur. J. Pharmacol.* 879: 173134.
- Li, Z., et al. 2020. N⁶-methyladenosine regulates glycolysis of cancer cells through PDK4. *Nat. Commun.* 11: 2578.
- Gao, Y., et al. 2020. IGF2BP3 and miR191-5p synergistically increase HCC cell invasiveness by altering ZO-1 expression. *Oncol. Lett.* 20: 1423-1431.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.