

SEN7 (G-7): sc-365794

BACKGROUND

SUMO (small ubiquitin-related modifier), a member of the ubiquitin-like protein family, regulates diverse cellular functions of a variety of target proteins, including transcription, DNA repair, nucleocytoplasmic trafficking and chromosome segregation. SUMO precursor proteins undergo cleavage of the residues after the "GG" region by SUMO-specific proteases in maturation. This cleavage of the precursor is a prerequisite for subsequent sumoylation. The sentrin-specific (or SUMO-specific) protease (SEN) proteins belong to the peptidase C48 family and include SENP1-3 and SENP5-8. SENP1, SENP2 and SENP3 degrade UBL1 and SMT3H2 conjugates and subsequently release the monomers from sumoylated substrates. HIPK2 is a desumoylation target for SENP1 which shuttles between the cytoplasm and the nucleus. Mutation analyses reveal that SENP1 contains the nuclear export sequence (NES) within the extreme carboxyl-terminal region, and SENP1 is exported to the cytoplasm in a NES-dependent manner. SENP2 has been implicated as a downregulator of CTNNB1 levels and may therefore be a modulator of the Wnt pathway. SUMO protease SENP3 reverses the sumoylation of MEF2 to augment its transcriptional and myogenic activities. SENP5 localizes to the nucleolus and preferentially processes SUMO-3. It is thought to play a role in mitosis and/or cytokinesis. SENP6 localizes to the cytoplasm and releases SUMO-1. Expression of SENP6 is higher in reproductive organs, indicating that it may mediate processes related to reproduction. SENP8 is involved in the release of sentrins.

REFERENCES

- Gong, L., et al. 2000. Differential regulation of sentrinized proteins by a novel sentrin-specific protease. *J. Biol. Chem.* 275: 3355-3359.
- Kim, K.I., et al. 2000. A new SUMO-1-specific protease, SUSP1, that is highly expressed in reproductive organs. *J. Biol. Chem.* 275: 14102-14106.
- Cheng, J., et al. 2004. SENP1 enhances androgen receptor-dependent transcription through desumoylation of histone deacetylase 1. *Mol. Cell Biol.* 24: 6021-6028.
- Reverter, D., et al. 2004. A basis for SUMO protease specificity provided by analysis of human SENP2 and a SENP2-SUMO complex. *Structure* 12: 1519-1531.
- Kim, Y.H., et al. 2005. Desumoylation of homeodomain-interacting protein kinase 2 (HIPK2) through the cytoplasmic-nuclear shuttling of the SUMO-specific protease SENP1. *FEBS Lett.* 579: 6272-6278.

CHROMOSOMAL LOCATION

Genetic locus: SENP7 (human) mapping to 3q12.3; Senp7 (mouse) mapping to 16 C1.1.

SOURCE

SEN7 (G-7) is a mouse monoclonal antibody raised against amino acids 912-956 mapping near the N-terminus of SENP7 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

SEN7 (G-7) is recommended for detection of SENP7 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for SENP7 siRNA (h): sc-61526, SENP7 siRNA (m): sc-61527, SENP7 shRNA Plasmid (h): sc-61526-SH, SENP7 shRNA Plasmid (m): sc-61527-SH, SENP7 shRNA (h) Lentiviral Particles: sc-61526-V and SENP7 shRNA (m) Lentiviral Particles: sc-61527-V.

Molecular Weight (predicted) of SENP7 isoforms: 120/116/28/101/112 kDa.


Molecular Weight (observed) of SENP7: 34 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, HUV-EC-C whole cell lysate: sc-364180 or PC-12 cell lysate: sc-2250.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



SEN7 (G-7): sc-365794. Western blot analysis of SENP7 expression in HUV-EC-C (A), HeLa (B), PC-12 (C), BC₃H1 (D) and TK-1 (E) whole cell lysates.



SEN7 (G-7): sc-365794. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human small intestine tissue showing cytoplasmic staining of glandular cells (B).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.