

Lsh (C-1): sc-365815

BACKGROUND

The SNF2 family of helicases are thought to act as transcriptional regulators by their ability to alter the structure of chromatin. One such member, lymphoid-specific helicase (Lsh, also designated Hells for lymphoid-specific DNA helicase), is highly expressed in lymphoid precursor cells in adult animals and is required for the proliferation of peripheral T lymphocytes. Lsh is also expressed in fetal liver and more abundantly in fetal thymus. Lsh protein shows substantial homology to other members of the SNF2 family that are involved in chromatin remodeling and transcription, however does not show similarity to members involved in DNA repair or recombination. The similarity of Lsh to another SNF2 homolog, Mi-2, which functions as a transcriptional silencer in chromatin remodeling, suggests that Lsh may participate in chromatin repression to regulate transcription, rather than chromatin "opening".

CHROMOSOMAL LOCATION

Genetic locus: HELLS (human) mapping to 10q23.33; Hells (mouse) mapping to 19 C3.

SOURCE

Lsh (C-1) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 259-287 near the N-terminus of Lsh of mouse origin.

PRODUCT

Each vial contains 200 µg IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-365815 X, 200 µg/0.1 ml.

Blocking peptide available for competition studies, sc-365815 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

Lsh (C-1) is recommended for detection of Lsh of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Lsh (C-1) is also recommended for detection of Lsh in additional species, including bovine.

Suitable for use as control antibody for Lsh siRNA (h): sc-38033, Lsh siRNA (m): sc-38034, Lsh shRNA Plasmid (h): sc-38033-SH, Lsh shRNA Plasmid (m): sc-38034-SH, Lsh shRNA (h) Lentiviral Particles: sc-38033-V and Lsh shRNA (m) Lentiviral Particles: sc-38034-V.

Lsh (C-1) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of Lsh: 100 kDa.

Positive Controls: JM1 whole cell lysate: sc-364233, CCRF-CEM cell lysate: sc-2225 or HeLa whole cell lysate: sc-2200.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



Lsh (C-1): sc-365815. Western blot analysis of Lsh expression in CCRF-CEM (A), Jurkat (B), HeLa (C) and NIH/3T3 (D) whole cell lysates.



Lsh (C-1): sc-365815. Western blot analysis of Lsh expression in CCRF-CEM (A) and JM1 (B) whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CONJUGATES

See **Lsh (H-4): sc-46665** for Lsh antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.