



ZAG (H-3): sc-365851

BACKGROUND

ZAG (Zn- α 2-glycoprotein, also designated Zn- α 2-gp) is a soluble, secreted protein found in serum and other body fluids (such as cerebrospinal fluid, blood plasma, urine and sweat). ZAG has a tendency to precipitate with zinc salts, has electrophoretic mobility in the region of the two globulins, and has 18% carbohydrate content. A member of the immunoglobulin superfamily, ZAG has a high degree of sequence similarity to class-I major histocompatibility complex (MHC) antigens. The ZAG structure includes a large groove analogous to MHC class I peptide binding grooves. The crystal structure of ZAG resembles a MHC class I heavy chain but does not bind the class I light chain β -2-Microglobulin, unlike other MHC related proteins. ZAG stimulates lipid degradation in adipocytes and its overexpression causes the extensive fat losses associated with some advanced cancers.

REFERENCES

1. Jirka, M. and Blanicky, P. 1973. Zn- α 2-glycoprotein in sweat. *Cas. Lek. Cesk.* 112: 1606-1608.
2. Ekman, R., Johansson, B.G. and Ravnskov, U. 1976. Renal handling of Zn- α 2-glycoprotein as compared with that of albumin and the retinol-binding protein. *J. Clin. Invest.* 57: 945-954.
3. Shibata, S. and Miura, K. 1982. Nephritogenic glycoprotein. IX. Plasma Zn- α 2-glycoprotein as a second source of nephritogenic glycoprotein in urine. *Nephron* 31: 170-176.
4. Uria, J.A., Fueyo, A., Balbin, M., Velasco, G., Pendas, A.M. and Lopez-Otin, C. 1996. Alternative splicing gives rise to two novel long isoforms of Zn- α 2-glycoprotein, a member of the immunoglobulin superfamily. *Gene* 169: 233-236.
5. Sanchez, L.M., Lopez-Otin, C. and Bjorkman, P.J. 1997. Biochemical characterization and crystalization of human Zn- α 2-glycoprotein, a soluble class I major histocompatibility complex homolog. *Proc. Natl. Acad. Sci. USA* 94: 4626-4630.
6. Sanchez, L.M., Chirino, A.J. and Bjorkman, P.J. 1999. Crystal structure of human ZAG, a fat-depleting factor related to MHC molecules. *Science* 283: 1914-1919.
7. Davidsson, P. and Nilsson, C.L. 1999. Peptide mapping of proteins in cerebrospinal fluid utilizing a rapid preparative two-dimensional electrophoretic procedure and matrix-assisted laser desorption/ionization mass spectrometry. *Biochim. Biophys. Acta* 1473: 391-399.

CHROMOSOMAL LOCATION

Genetic locus: AZGP1 (human) mapping to 7q22.1.

SOURCE

ZAG (H-3) is a mouse monoclonal antibody raised against amino acids 13-135 mapping near the N-terminus of ZAG of human origin.

PRODUCT

Each vial contains 200 μ g IgG γ 1 kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

ZAG (H-3) is recommended for detection of ZAG of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for ZAG siRNA (h): sc-36865, ZAG shRNA Plasmid (h): sc-36865-SH and ZAG shRNA (h) Lentiviral Particles: sc-36865-V.


Molecular Weight of ZAG: 47 kDa.

Positive Controls: human spleen extract: sc-363779 or human plasma extract: sc-364374..


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



ZAG (H-3): sc-365851. Western blot analysis of ZAG in human plasma.



ZAG (H-3): sc-365851. Western blot analysis of ZAG expression in human spleen tissue extract.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.