SANTA CRUZ BIOTECHNOLOGY, INC.

GSS (C-5): sc-365863



BACKGROUND

GSS (glutathione synthetase) is a 474 amino acid protein encoded by the gene located at human chromosome 20g11.22. GSS consists of three loops projecting from an antiparallel β -sheet, a parallel β -sheet and a lid of antiparallel sheets, which provide access to the ATP-binding site. Although Southern blot and gene analysis suggest that GSS may be the only member of a unique family, the crystal structure indicates that GSS belongs to the ATP-GRASP superfamily. GSS is expressed in hemocytes and nucleated cells, including the brain. GSS occurs as a homodimer. There are two steps in the production of glutathione, begining with γ -GCS and ending with GSS. In an ATP-dependent reaction, GSS produces glutathione from y-glutamylcysteine and glycine precursors. Partial hepatectomy, diethyl maleate, buthionine sulfoximine, tert-butylhaydroguinone and thioacetamide increase the ex-pression of GSS, which causes an increase in glutathione levels. An inherited autosomal recessive disorder, 5-oxoprolinuria (pyroglutamic aciduria), is caused by GSS deficiencies, which leads to central nervous system damage, hemolytic anemia, metabolic acidosis and urinary excretion of 5-oxoproline. A missense mutation in the gene encoding GSS leads to a GSS deficiency restricted to erythrocytes, which causes only hemolytic anemia.

REFERENCES

- 1. Webb, G.C., et al. 1995. The gene encoding human glutathione synthetase (GSS) maps to the long arm of chromosome 20 at band 11.2. Genomics 30: 617-619.
- Gali, R.R. and Board, P.G. 1995. Sequencing and expression of a cDNA for human glutathione synthetase. Biochem. J. 310: 353-358
- Shi, Z.Z., et al. 1996. Mutations in the glutathione synthetase gene cause 5-oxoprolinuria. Nat. Genet. 14: 361-365.
- Polekhina, G., et al. 1999. Molecular basis of glutathione synthetase deficiency and a rare gene permutation event. EMBO J. 18: 3204-3213.

CHROMOSOMAL LOCATION

Genetic locus: GSS (human) mapping to 20q11.22; Gss (mouse) mapping to 2 H1.

SOURCE

GSS (C-5) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 357-385 within an internal region of GSS of human origin.

PRODUCT

Each vial contains 200 μg lgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-365863 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

STORAGE

Store at 4° C, **D0 NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

GSS (C-5) is recommended for detection of GSS of mouse, rat, human, and, to a lesser extent, hamster origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffinembedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

GSS (C-5) is also recommended for detection of GSS in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for GSS siRNA (h): sc-41980, GSS siRNA (m): sc-41981, GSS shRNA Plasmid (h): sc-41980-SH, GSS shRNA Plasmid (m): sc-41981-SH, GSS shRNA (h) Lentiviral Particles: sc-41980-V and GSS shRNA (m) Lentiviral Particles: sc-41981-V.

Molecular Weight of GSS: 52 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, COLO 205 whole cell lysate: sc-364177 or mouse brain extract: sc-2253.

DATA





GSS (C-5): sc-365863. Western blot analysis of GSS expression in HeLa (A), COLO 205 (B), Hep G2 (C) and EOC 20 (D) whole cell lysates and mouse brain tissue extract (E).

GSS (C-5): sc-365863. Immunoperoxidase staining of formalin fixed, paraffin-embedded human uterine cervix tissue showing cytoplasmic staining of squamous epithelial cells.

SELECT PRODUCT CITATIONS

- Yin, Z.X., et al. 2017. PARP-1 inhibitors sensitize HNSCC cells to APR-246 by inactivation of thioredoxin reductase 1 (TrxR1) and promotion of Ros accumulation. Oncotarget 9: 1885-1897.
- Li, S., et al. 2019. Glutathione contributes to efficient post-Golgi trafficking of incoming HPV16 genome. PLoS ONE 14: e0225496.
- 3. Jara, O., et al. 2020. Do connexin mutants cause cataracts by perturbing glutathione levels and redox metabolism in the lens? Biomolecules 10: 1418.
- Patten, D.A., et al. 2021. Altered mitochondrial fusion drives defensive glutathione synthesis in cells able to switch to glycolytic ATP production. Biochim. Biophys. Acta Mol. Cell Res. 1868: 118854.

RESEARCH USE

For research use only, not for use in diagnostic procedures.