

BACE (A-12): sc-365948

BACKGROUND

Autosomal dominant Alzheimer's disease is caused by mutations in the gene encoding the β -Amyloid protein precursor (APP). Amyloid β -peptide (A β), the major feature of amyloid plaques in Alzheimer's patients, is the product of APP cleavage by β - and γ -secretases. BACE is the transmembrane protease which cleaves A β from APP. BACE and the related protein Asp1 are both widely expressed in human tissue with the highest levels in the pancreas. BACE is localized within Golgi and endosomes.

REFERENCES

1. Kang, J., et al. 1987. The precursor of Alzheimer's disease Amyloid A4 protein resembles a cell-surface receptor. *Nature* 325: 733-736.
2. Goate, A., et al. 1991. Segregation of a missense mutation in the amyloid precursor protein gene with familial Alzheimer's disease. *Nature* 349: 704-706.
3. Mullan, M., et al. 1992. A pathogenic mutation for probable Alzheimer's disease in the APP gene at the N-terminus of β -Amyloid. *Nat. Genet.* 1: 345-347.
4. Selkoe, D.J. 1998. The cell biology of β -Amyloid precursor protein and presenilin in Alzheimer's disease. *Trends Cell Biol.* 8: 447-453.

CHROMOSOMAL LOCATION

Genetic locus: BACE1 (human) mapping to 11q23.3; Bace1 (mouse) mapping to 9 A5.2.

SOURCE

BACE (A-12) is a mouse monoclonal antibody raised against amino acids 419-501 of BACE of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

BACE (A-12) is recommended for detection of BACE of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for BACE siRNA (h): sc-37224, BACE siRNA (m): sc-37225, BACE shRNA Plasmid (h): sc-37224-SH, BACE shRNA Plasmid (m): sc-37225-SH, BACE shRNA (h) Lentiviral Particles: sc-37224-V and BACE shRNA (m) Lentiviral Particles: sc-37225-V.


Molecular Weight of BACE: 70 kDa.

Positive Controls: SH-SY5Y cell lysate: sc-3812 or BACE (h): 293T Lysate: sc-159912.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



BACE (A-12): sc-365948. Western blot analysis of BACE expression in non-transfected: sc-117752 (A) and human BACE transfected: sc-159912 (B) 293T whole cell lysates.



BACE (A-12): sc-365948. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human colon tissue showing cytoplasmic staining of glandular cells (B).

SELECT PRODUCT CITATIONS

1. Chen, L., et al. 2016. Neuroprotective effects of vitexin against isoflurane-induced neurotoxicity by targeting the TRPV1 and NR2B signaling pathways. *Mol. Med. Rep.* 14: 5607-5613.
2. Li, W.H., et al. 2021. Deletion of Dcf1 reduces Amyloid- β aggregation and mitigates memory deficits. *J. Alzheimers Dis.* 81: 1181-1194.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **BACE (61-3E7): sc-33711** for BACE antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.