

Misato (H-76): sc-366025

BACKGROUND

Drosophila melanogaster is a proven and effective model for studying developmental and cellular processes common to higher eukaryotes. Approximately 13,600 genes have been elucidated from more than 120 megabases of euchromatin, and they are organized among the chromosomes 2, 3, 4, X and Y, with the Y chromosome being predominately heterochromatic. *Drosophila* genes can be categorized based on the type of protein for which they encode and are represented by six major classifications, which include intracellular signaling proteins, transmembrane proteins, RNA binding proteins, secreted factors, transcription regulators (basic helix-loop-helix, homeodomain containing, zinc finger containing, and chromatin associated) or other functional proteins. The misato gene encodes a protein that contains a mixture of peptide motifs found in α -, β -, and γ -tubulins, as well as a motif related to part of the myosin heavy chain proteins. Null mutations at the misato locus of *Drosophila melanogaster* are associated with irregular chromosomal segregation at cell division and result in larvae that have reduced levels of imaginal disk tissue, a reduction in brain size, and die during the larval stage of development.

REFERENCES

- Adams, M.D., Celniker, S.E., Holt, R.A., Evans, C.A., Gocayne, J.D., Amanatides, P.G., Scherer, S.E., Li, P.W., Hoskins, R.A., Galle, R.F., Georg, R.A., Lewis, S.E., Richards, S., Ashburner, M., et al. 2000. The genome sequence of *Drosophila melanogaster*. *Science* 287: 2185-2195.
- Miklos, G.L., Yamamoto, M., Burns, R.G. and Maleszka, R. 1997. An essential cell division gene of *Drosophila*, absent from *Saccharomyces*, encodes an unusual protein with tubulin-like and myosin-like peptide motifs. *Proc. Natl. Acad. Sci. USA* 94: 5189-5194.
- Nogales, E., Wolf, S.G. and Downing, K.H. 1998. Structure of the α β tubulin dimer by electron crystallography. *Nature* 391: 199-203.
- LocusLink Report (LocusID: 33119). <http://www.ncbi.nlm.nih.gov/LocusLink/>

CHROMOSOMAL LOCATION

Genetic locus: MSTO1 (human) mapping to 1q22; Msto1 (mouse) mapping to 3 F1.

SOURCE

Misato (H-76) is a rabbit polyclonal antibody raised against amino acids 1-76 mapping at the N-terminus of Misato of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

Misato (H-76) is recommended for detection of Misato of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Misato (H-76) is also recommended for detection of Misato in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Misato siRNA (h): sc-62619, Misato siRNA (m): sc-62620, Misato shRNA Plasmid (h): sc-62619-SH, Misato shRNA Plasmid (m): sc-62620-SH, Misato shRNA (h) Lentiviral Particles: sc-62619-V and Misato shRNA (m) Lentiviral Particles: sc-62620-V.

Molecular Weight of Misato: 62 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.