

Tak1 siRNA (m): sc-36607

BACKGROUND

Several serine/threonine protein kinases have been implicated as intermediates in signal transduction pathways. These include ERK/MAP kinases, ribosomal S6 kinase (Rsk) and Raf-1. Raf-1 has intrinsic kinase activity towards serine/threonine residues and that is widely expressed in many tissue types and cell lines. Raf-1 activation is dependent on the small molecular weight GTPase Ras, but the means by which this activation occurs is poorly understood. Two proteins putatively involved in this process are Ksr-1 and Tak1. Ksr-1 (kinase suppressor of Ras) is a novel Raf-related protein kinase whose function is required for Ras signal transduction. Whether Ksr-1 lies directly downstream of Ras or acts in a parallel pathway is not yet known. Tak1 (TGF β -activated kinase) has been shown to participate in the activation of the MAP kinase family in response to TGF β stimulation.

CHROMOSOMAL LOCATION

Genetic locus: Map3k7 (mouse) mapping to 4 A5.

PRODUCT

Tak1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Tak1 shRNA Plasmid (m): sc-36607-SH and Tak1 shRNA (m) Lentiviral Particles: sc-36607-V as alternate gene silencing products.

For independent verification of Tak1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-36607A, sc-36607B and sc-36607C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Tak1 siRNA (m) is recommended for the inhibition of Tak1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Tak1 (C-9): sc-7967 is recommended as a control antibody for monitoring of Tak1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Tak1 gene expression knockdown using RT-PCR Primer: Tak1 (m)-PR: sc-36607-PR (20 μ l, 458 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

- Haase, R., et al. 2005. Yersinia outer protein P suppresses TGF- β -activated kinase-1 activity to impair innate immune signaling in *Yersinia enterocolitica*-infected cells. *J. Immunol.* 175: 8209-8217.
- Yang, Y., et al. 2007. NOD2 pathway activation by MDP or *Mycobacterium tuberculosis* infection involves the stable polyubiquitination of Rip2. *J. Biol. Chem.* 282: 36223-36229.
- Erdogan, M., et al. 2008. Transforming growth factor- β (TGF- β) and TGF- β -associated kinase 1 are required for R-Ras-mediated transformation of mammary epithelial cells. *Cancer Res.* 68: 6224-6231.
- Villar, A.V., et al. 2013. BAMBI (BMP and activin membrane-bound inhibitor) protects the murine heart from pressure-overload biomechanical stress by restraining TGF- β signaling. *Biochim. Biophys. Acta* 1832: 325-335.
- Song, Z., et al. 2014. Roles of the kinase Tak1 in CD40-mediated effects on vascular oxidative stress and neointima formation after vascular injury. *PLoS ONE* 9: e101671.
- Campo, G.M., et al. 2015. β -Arrestin 1 is involved in the catabolic response stimulated by hyaluronan degradation in mouse chondrocytes. *Cell Tissue Res.* 361: 567-579.
- Chen, H., et al. 2016. PGC-1 β suppresses saturated fatty acid-induced macrophage inflammation by inhibiting Tak1 activation. *IUBMB Life* 68: 145-155.
- Nakajima, M., et al. 2017. IL-17F induces IL-6 via Tak1-NF κ B pathway in airway smooth muscle cells. *Immun. Inflamm. Dis.* 5: 124-131.
- Zhou, J., et al. 2018. Tak1 mediates excessive autophagy via p38 and ERK in cisplatin-induced acute kidney injury. *J. Cell. Mol. Med.* 22: 2908-2921.
- Liu, M., et al. 2020. Macrophage K63-linked ubiquitination of YAP promotes its nuclear localization and exacerbates atherosclerosis. *Cell Rep.* 32: 107990.
- Tenshin, H., et al. 2022. TGF- β -activated kinase-1 inhibitor LL-Z1640-2 reduces joint inflammation and bone destruction in mouse models of rheumatoid arthritis by inhibiting NLRP3 inflammasome, TACE, TNF- α and RANKL expression. *Clin. Transl. Immunology* 11: e1371.

RESEARCH USE

For research use only, not for use in diagnostic procedures.