

KCC1 (H-69): sc-366076

BACKGROUND

The four isoforms of potassium/chloride co-transport channels (KCC) belong to a superfamily of cation-chloride co-transporters involved in cell volume maintenance. Nitric oxide (NO) donors activate KCCs, while inhibitors of the cGMP pathway prevent NO donor activation. The ubiquitously expressed KCC1 contains 12 transmembrane domains with both cytoplasmic N and C terminal domains. KCC2 expression is limited to neuronal tissues by a restrictive element similar to the neuronal-restrictive silencing factor. In neurons, KCC2 expression is correlated with an inhibitory response to GABA, while the absence of KCC2 is necessary for an unusual excitatory response to GABA. Alterations of KCC2 expression in the inferior colliculus of rat brain may be related to seizure susceptibility. Conversely, KCC3 is not suspected to play a major role in epilepsy. The two splice variants of KCC3, KCC3a and KCC3b, are predominantly expressed in brain and kidney, respectively, while KCC4 is expressed in muscle, brain, lung, heart and kidney.

REFERENCES

1. Mount, D.B., et al. 1999. Cloning and characterization of KCC3 and KCC4, new members of the cation-chloride cotransporter gene family. *J. Biol. Chem.* 274: 16355-16362.
2. Lauf, P.K., et al. 2000. K-Cl cotransport: properties and molecular mechanism. *Cell. Physiol. Biochem.* 10: 341-354.
3. Di Fulvio, M., et al. 2001. Protein kinase G regulates potassium chloride cotransporter-3 expression in primary cultures of rat vascular smooth muscle cells. *J. Biol. Chem.* 276: 21046-21052.
4. Lauf, P.K., et al. 2001. K-Cl cotransport: immunohistochemical and ion flux studies in human embryonic kidney (HEK293) cells transfected with full-length and C-terminal-domain-truncated KCC1 cDNAs. *Cell. Physiol. Biochem.* 11: 143-160.
5. Karadsheh, M.F., et al. 2001. Neuronal restrictive silencing element is found in the KCC2 gene: molecular basis for KCC2-specific expression in neurons. *J. Neurophysiol.* 85: 995-997.
6. Hubner, C.A., et al. 2001. Disruption of KCC2 reveals an essential role of K-Cl cotransport already in early synaptic inhibition. *Neuron* 30: 515-524.
7. Kanaka, C., et al. 2001. The differential expression patterns of messenger RNAs encoding K-Cl cotransporters (KCC1,2) and Na-K-Cl cotransporter (NKCC1) in the rat nervous system. *Neuroscience* 104: 933-946.
8. Reid, K.H., et al. 2001. The mRNA level of the potassium-chloride cotransporter KCC2 covaries with seizure susceptibility in inferior colliculus of the post-ischemic audiogenic seizure-prone rat. *Neurosoci. Lett.* 308: 29-32.
9. Steinlein, O.K., et al. 2001. Mutation analysis of the potassium chloride cotransporter KCC3 (SLC12A6) in rolandic and idiopathic generalized epilepsy. *Epilepsy Res.* 44: 191-195.

CHROMOSOMAL LOCATION

Genetic locus: SLC12A4 (human) mapping to 16q22.1; Slc12a4 (mouse) mapping to 8 D3.

SOURCE

KCC1 (H-69) is a rabbit polyclonal antibody raised against amino acids 1-69 mapping at the N-terminus of KCC1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

KCC1 (H-69) is recommended for detection of KCC1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

KCC1 (H-69) is also recommended for detection of KCC1 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for KCC1 siRNA (h): sc-42604, KCC1 siRNA (m): sc-42605, KCC1 shRNA Plasmid (h): sc-42604-SH, KCC1 shRNA Plasmid (m): sc-42605-SH, KCC1 shRNA (h) Lentiviral Particles: sc-42604-V and KCC1 shRNA (m) Lentiviral Particles: sc-42605-V.

Molecular Weight of KCC1: 121/119/112/85 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **KCC1 (FT-94): sc-134370**, our highly recommended monoclonal alternative to KCC1 (H-69).