

# PI-9 (M-244): sc-366114

## BACKGROUND

Serine proteinase inhibitors (serpins) function as regulators of serine proteinase activity in a variety of physiological processes. Proteinase inhibitor-9 (PI-9, also designated cytoplasmic antiproteinase 3, or CAP3) is a member of the ovalbumin family of serpins that is expressed in placenta, lung and cytotoxic lymphocytes. PI-9 is a potent inhibitor of granzyme B and of granzyme B-mediated apoptosis, and is also an inhibitor of caspase-1 and, to a lesser extent, caspase-4 and caspase-8. Because granzyme B promotes DNA degradation and rapidly translocates to the nucleus to bind to a nuclear component, PI-9 is present in the nuclei of human cytotoxic cells, endothelial cells and epithelial cells. PI-9 is exported from nuclei via a leptomycin B-sensitive pathway, suggesting that the nucleocytoplasmic distribution of PI-9 involves a non-conventional nuclear import pathway and the export factor CRM1. Estrogen rapidly and strongly induces PI-9, which is an estrogen-regulated human gene. PI-9 expression is also upregulated in response to inflammatory stimuli. This upregulation protects cells from apoptosis induced by endogenously expressed or released granzyme B, particularly during target cell killing. In addition, PI-9 is expressed in a variety of human and murine tumors.

## REFERENCES

- Dahlen, J.R., Foster, D.C. and Kisiel, W. 1997. Human proteinase inhibitor 9 (PI9) is a potent inhibitor of subtilisin A. *Biochem. Biophys. Res. Commun.* 238: 329-333.
- Sun, J., Ooms, L., Bird, C.H., Sutton, V.R., Trapani, J.A. and Bird, P.I. 1997. A new family of 10 murine ovalbumin serpins includes two homologs of proteinase inhibitor 8 and two homologs of the granzyme B inhibitor (proteinase inhibitor 9). *J. Biol. Chem.* 272: 15434-15441.
- Sun, J., Stephens, R., Mirza, G., Kanai, H., Ragoussis, J. and Bird, P.I. 1998. A serpin gene cluster on human chromosome 6p25 contains PI6, PI9 and dELAN<sub>H2</sub> which have a common structure almost identical to the 18q21 ovalbumin serpin genes. *Cytogenet. Cell Genet.* 82: 273-277.
- Dahlen, J.R., Foster, D.C. and Kisiel, W. 1999. Inhibition of neutrophil elastase by recombinant human proteinase inhibitor 9. *Biochem. Biophys. Acta* 1451: 233-241.
- Annand, R.R., Dahlen, J.R., Sprecher, C.A., De Dreu, P., Foster, D.C., Mankovich, J.A., Talanian, R.V., Kisiel, W. and Giegel, D.A. 1999. Caspase-1 (interleukin-1 $\beta$ -converting enzyme) is inhibited by the human serpin analog proteinase inhibitor 9. *Biochem. J.* 342: 655-665.
- Kanamori, H., Kreig, S., Mao, C., Di Pippo, V.A., Wang, S., Zajchowski, D.A. and Shapiro, D.J. 2000. Proteinase inhibitor 9, an inhibitor of granzyme B-mediated apoptosis, is a primary estrogen-inducible gene in human liver cells. *J. Biol. Chem.* 275: 5867-5873.
- Bird, C.H., Blink, E.J., Hirst, C.E., Buzza, M.S., Steele, P.M., Sun, J., Jans, D.A. and Bird, P.I. 2001. Nucleocytoplasmic distribution of the ovalbumin serpin PI-9 requires a nonconventional nuclear import pathway and the export factor Crm1. *Mol. Cell. Biol.* 21: 5396-5407.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## CHROMOSOMAL LOCATION

Genetic locus: Serpinb9 (mouse) mapping to 13 A3.3.

## SOURCE

PI-9 (M-244) is a rabbit polyclonal antibody raised against amino acids 77-320 mapping within an internal region of PI-9 of mouse origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

PI-9 (M-244) is recommended for detection of PI-9 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for PI-9 siRNA (m): sc-152245, PI-9 shRNA Plasmid (m): sc-152245-SH and PI-9 shRNA (h) Lentiviral Particles: sc-40949-V.

Molecular Weight of PI-9: 42 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker<sup>™</sup> compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz<sup>™</sup> Mounting Medium: sc-24941.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.


 MONOS  
 Satisfation  
 Guaranteed

Try **PI-9 (F-6): sc-390501** or **PI-9 (F-12): sc-390561**, our highly recommended monoclonal alternatives to PI-9 (M-244).