

IL-27 (H-59): sc-366222

BACKGROUND

IL-27 triggers expansion of antigen-specific naive CD4⁺ T cells and promotes polarization towards a Th1 phenotype with expression of γ -interferon. IL-27 contributes to the development of an adaptive immune response through its action on CD4⁺ T cells and also directly acts on cells of the innate immune system. IL-27 protein levels increase upon activation of antigen-presenting cells. IL-27 protein induces orphan cytokine receptor WSX-1/TCCR-dependent clonal expansion of naive but not memory CD4⁺ T cells. IL-27 signaling through TCCR/WSX-1 induces phosphorylation of Stat1-5. The predicted 243 amino acid human IL-27 protein, which is 73% identical to the mouse protein, contains an N-terminal signal peptide, several O-glycosylation sites and a stretch of 13 glutamate residues between helices C and D.

REFERENCES

- Pflanz, S., et al. 2002. IL-27, a heterodimeric cytokine composed of EBI3 and p28 protein, induces proliferation of naive CD4⁺ T cells. *Immunity* 16: 779-790.
- Cordoba-Rodriguez, R., et al. 2003. L-23 and IL-27: new members of the growing family of IL-12-related cytokines with important implications for therapeutics. *Exp. Opin. Biol. Ther.* 3: 715-723.
- Lucas, S., et al. 2003. IL-27 regulates IL-12 responsiveness of naive CD4⁺ T cells through Stat1-dependent and -independent mechanisms. *Proc. Natl. Acad. Sci. USA* 100: 15047-15052.
- Villarino, A.V., et al. 2004. Understanding the pro- and anti-inflammatory properties of IL-27. *J. Immunol.* 173: 715-720.
- Goldberg, R., et al. 2004. Suppression of ongoing adjuvant-induced arthritis by neutralizing the function of the p28 subunit of IL-27. *J. Immunol.* 173: 1171-1178.
- Yoshimoto, T., et al. 2004. Induction of IgG_{2a} class switching in B cells by IL-27. *J. Immunol.* 173: 2479-2485.
- Artis, D., et al. 2004. The IL-27 receptor (WSX-1) is an inhibitor of innate and adaptive elements of type 2 immunity. *J. Immunol.* 173: 5626-5634.
- LocusLink Report (LocusID: 53342). <http://www.ncbi.nlm.nih.gov/LocusLink/>

CHROMOSOMAL LOCATION

Genetic locus: IL17D (human) mapping to 13q12.11.

SOURCE

IL-27 (H-59) is a rabbit polyclonal antibody raised against amino acids 60-118 mapping within an internal region of IL-27 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

IL-27 (H-59) is recommended for detection of IL-27 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for IL-27 siRNA (h): sc-72184, IL-27 shRNA Plasmid (h): sc-72184-SH and IL-27 shRNA (h) Lentiviral Particles: sc-72184-V.

Molecular Weight of IL-27: 27 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **IL-27 (LB-L22): sc-134367**, our highly recommended monoclonal alternative to IL-27 (H-59).