

TNF-R2 siRNA (m): sc-36690

BACKGROUND

Tumor necrosis factor (TNF) is a pleiotropic cytokine whose function is mediated through two distinct cell surface receptors. These receptors, designated TNF-R1 and TNF-R2, are expressed on most cell types. The majority of TNF functions are primarily mediated through TNF-R1, while signaling through TNF-R2 occurs less extensively and is confined to cells of the immune system. Both of these proteins belong to the growing TNF and nerve growth factor (NGF) receptor superfamily, which includes FAS, CD30, CD27 and CD40. The members of this superfamily are type I membrane proteins that share sequence homology confined to the extracellular region. TNF-R1 shares a motif termed the "death domain" with FAS and three structurally unrelated signaling proteins, TRADD, FADD and RIP. This death domain is required for transduction of the apoptotic signal.

REFERENCES

1. Smith, C.A., et al. 1994. The TNF receptor superfamily of cellular and viral proteins: activation, costimulation, and death. *Cell* 76: 959-962.
2. Nagata, S. and Golstein, P. 1995. The FAS death factor. *Science* 267: 1449-1456.
3. Cleveland, J.L. and Ihle, J.N. 1995. Contenders in FAS-L/TNF death signaling. *Cell* 81: 479-482.
4. Hsu, H., et al. 1995. The TNF receptor 1-associated protein TRADD signals cell death and NF κ B activation. *Cell* 81: 495-504.
5. Chinnaiyan, A.M., et al. 1995. FADD, a novel death domain-containing protein, interacts with the death domain of FAS and initiates apoptosis. *Cell* 81: 505-512.
6. Stanger, B.Z., et al. 1995. RIP: a novel protein containing a death domain that interacts with FAS/Apo-1 (CD95) in yeast and causes cell death. *Cell* 81: 513-523.
7. Boldin, M.P., et al. 1995. Self-association of the "death domains" of the p55 tumor necrosis factor (TNF) receptor and FAS/Apo-1 prompts signaling for TNF and FAS/Apo-1 effects. *J. Biol. Chem.* 270: 387-391.
8. Hofmann, K. and Tschoop, J. 1995. The death domain motif found in FAS (Apo-1) and TNF receptor is present in proteins involved in apoptosis and axonal guidance. *FEBS Lett.* 371: 321-323.

CHROMOSOMAL LOCATION

Genetic locus: Tnfrsf1b (mouse) mapping to 4 E1.

PRODUCT

TNF-R2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TNF-R2 shRNA Plasmid (m): sc-36690-SH and TNF-R2 shRNA (m) Lentiviral Particles: sc-36690-V as alternate gene silencing products.

For independent verification of TNF-R2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-36690A, sc-36690B and sc-36690C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TNF-R2 siRNA (m) is recommended for the inhibition of TNF-R2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

TNF-R2 (D-2): sc-8041 is recommended as a control antibody for monitoring of TNF-R2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TNF-R2 gene expression knockdown using RT-PCR Primer: TNF-R2 (m)-PR: sc-36690-PR (20 μ l, 431 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Song, L., et al. 2007. p85 α acts as a novel signal transducer for mediation of cellular apoptotic response to UV radiation. *Mol. Cell. Biol.* 27: 2713-2731.
2. Li, H., et al. 2015. Administration of progranulin (PGRN) triggers ER stress and impairs insulin sensitivity via PERK-eIF2 α -dependent manner. *Cell Cycle* 14: 1893-1907.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.