

TRAF6 siRNA (h): sc-36717

BACKGROUND

Tumor necrosis factor receptor-associated factor 6 (TRAF6) regulates adaptive immunity, innate immunity and bone metabolism. TRAF6 is a ubiquitin (Ub) ligase that mediates the activation of protein kinases, such as transforming growth factor β -activated kinase (TAK1) and I κ B kinase (IKK), by catalyzing the formation of a unique polyubiquitin chain linked through Lys 63 of Ub. TRAF6 is essential for activating NF κ B signaling pathway in response to interleukin-1 and Toll-like receptor ligands. The coiled-coil domain of TRAF6 is essential for its auto-ubiquitination and activating NF κ B signaling pathway. TRAF6 interacts with various protein kinases including IRAK1/IRAK, SRC and PKC ζ , which provides a link between distinct signaling pathways.

CHROMOSOMAL LOCATION

Genetic locus: TRAF6 (human) mapping to 11p12.

PRODUCT

TRAF6 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TRAF6 shRNA Plasmid (h): sc-36717-SH and TRAF6 shRNA (h) Lentiviral Particles: sc-36717-V as alternate gene silencing products.

For independent verification of TRAF6 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-36717A, sc-36717B and sc-36717C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TRAF6 siRNA (h) is recommended for the inhibition of TRAF6 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

TRAF6 (D-10): sc-8409 is recommended as a control antibody for monitoring of TRAF6 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TRAF6 gene expression knockdown using RT-PCR Primer: TRAF6 (h)-PR: sc-36717-PR (20 μ l, 510 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

- Guasparri, I., et al. 2006. The KSHV oncoprotein vFLIP contains a TRAF-interacting motif and requires TRAF2 and TRAF3 for signalling. *EMBO Rep.* 7: 114-119.
- Guasparri, I., et al. 2008. EBV LMP2A affects LMP1-mediated NF κ B signaling and survival of lymphoma cells by regulating TRAF2 expression. *Blood* 111: 3813-3820.
- Ahmad, R., et al. 2008. Requirement of TLR2-mediated signaling for the induction of IL-15 gene expression in human monocytic cells by HSV-1. *Blood* 112: 2360-2368.
- Ahmad, R., et al. 2009. Adaptor proteins and Ras synergistically regulate IL-1-induced ADAMTS-4 expression in human chondrocytes. *J. Immunol.* 182: 5081-5087.
- Tang, L., et al. 2011. Expression of TRAF6 and pro-inflammatory cytokines through activation of TLR2, TLR4, NOD1, and NOD2 in human periodontal ligament fibroblasts. *Arch. Oral Biol.* 56: 1064-1072.
- Paik, J.H., et al. 2011. MicroRNA-146a downregulates NF κ B activity via targeting TRAF6 and functions as a tumor suppressor having strong prognostic implications in NK/T cell lymphoma. *Clin. Cancer Res.* 17: 4761-4771.
- Li, L., et al. 2012. Viral oncoprotein LMP1 disrupts p53-induced cell cycle arrest and apoptosis through modulating K63-linked ubiquitination of p53. *Cell Cycle* 11: 2327-2336.
- Lin, D., et al. 2014. IL-17 regulates the expressions of RANKL and OPG in human periodontal ligament cells via TRAF6/TBK1-JNK/NF κ B pathways. *Immunology* 144: 472-485.
- Chan, S.T., et al. 2016. Suppression of host innate immune response by hepatitis C virus via induction of autophagic degradation of TRAF6. *J. Virol.* 90: 10928-10935.
- Wang, X.L., et al. 2016. Rosuvastatin attenuates CD40L-induced down-regulation of extracellular matrix production in human aortic smooth muscle cells via TRAF6-JNK-NF κ B pathway. *PLoS ONE* 11: e0153919.
- Li, L., et al. 2018. RIG-I is involved in inflammation through the IPS-1/TRAF6 pathway in astrocytes under chemical hypoxia. *Neurosci. Lett.* 672: 46-52.

RESEARCH USE

For research use only, not for use in diagnostic procedures.