

LAT (B-3): sc-373706

BACKGROUND

T cell receptors activate immune responses by recognizing antigen and initiating a cascade of intracellular signal transduction events, eventually culminating in cell proliferation and differentiation. Both protein tyrosine kinases and PLC γ are activated by this event. LAT, or linker for activation of T cells, is an integral membrane protein that has been shown to associate with PLC γ 1, as well as GRB2 and the p85 subunit of PI 3-kinase. Binding of these signaling molecules to LAT is associated with phosphorylation of LAT by ZAP-70/Syk tyrosine kinases. LAT appears to play a role in activation of transcription mediated by AP-1 and NFAT following stimulation of the T cell receptor, suggesting that it acts as a linker protein in T cell activation. LAT protein is palmitoylated, and this modification is required for its tyrosine phosphorylation and localization to glycolipid-enriched microdomains.

REFERENCES

- Weiss, A., et al. 1991. Signal transduction by the T cell antigen receptor. *Semin. Immunol.* 3: 313-324.
- Isakov, N., et al. 1994. The role of tyrosine kinases and phosphotyrosine-containing recognition motifs in regulation of the T cell-antigen receptor-mediated signal transduction pathway. *J. Leukoc. Biol.* 55: 265-271.
- Zhang, W., et al. 1998. LAT: the ZAP-70 tyrosine kinase substrate that links T cell receptor to cellular activation. *Cell* 92: 83-92.
- Cantrell, D. 1998. The real LAT steps forward. *Trends Cell Biol.* 8: 180-182.
- Zhang, W., et al. 1998. LAT palmitoylation: its essential role in membrane microdomain targeting and tyrosine phosphorylation during T cell activation. *Immunity* 9: 239-246.
- Brdicka, T., et al. 1998. T cell receptor signalling results in rapid tyrosine phosphorylation of the linker protein LAT present in detergent-resistant membrane microdomains. *Biochem. Biophys. Res. Commun.* 248: 356-360.

CHROMOSOMAL LOCATION

Genetic locus: LAT (human) mapping to 16p11.2; Lat (mouse) mapping to 7 F3.

SOURCE

LAT (B-3) is a mouse monoclonal antibody raised against amino acids 1-233 representing full length LAT of human origin.

PRODUCT

Each vial contains 200 μ g IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

LAT (B-3) is available conjugated to agarose (sc-373706 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-373706 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-373706 PE), fluorescein (sc-373706 FITC), Alexa Fluor[®] 488 (sc-373706 AF488), Alexa Fluor[®] 546 (sc-373706 AF546), Alexa Fluor[®] 594 (sc-373706 AF594) or Alexa Fluor[®] 647 (sc-373706 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-373706 AF680) or Alexa Fluor[®] 790 (sc-373706 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

LAT (B-3) is recommended for detection of LAT of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for LAT siRNA (h): sc-35795, LAT siRNA (m): sc-35796, LAT shRNA Plasmid (h): sc-35795-SH, LAT shRNA Plasmid (m): sc-35796-SH, LAT shRNA (h) Lentiviral Particles: sc-35795-V and LAT shRNA (m) Lentiviral Particles: sc-35796-V.


Molecular Weight of LAT: 36-38 kDa.

Positive Controls: LAT (m): 293T Lysate: sc-127084, HuT 78 whole cell lysate: sc-2208 or BYDP whole cell lysate: sc-364368.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA



LAT (B-3): sc-373706. Western blot analysis of LAT expression in non-transfected 293T: sc-117752 (A), mouse LAT transfected 293T: sc-127084 (B), BYDP (C) and HuT 78 (D) whole cell lysates.



LAT (B-3): sc-373706. Western blot analysis of LAT expression in WR19L whole cell lysate (A) and rat thymus tissue extract (B).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.