SANTA CRUZ BIOTECHNOLOGY, INC.

E-FABP (D-12): sc-373788



BACKGROUND

Fatty acid-binding proteins, designated FABPs, are a family of homologous, 14-15 kDa cytoplasmic proteins that are expressed in a highly tissue-specific manner and play an integral role in the balance between lipid and carbohydrate metabolism. FABPs mediate fatty acid (FA) and/or hydrophobic ligand uptake, transport, and targeting within their respective tissues. The mechanisms underlying these actions can give rise to both passive diffusional uptake and protein-mediated transmembrane transport of FAs. FABPs are expressed in adipocytes (A-FABP), brain, (B-FABP), epithelium (E-FABP, psoriasis-associated FABP, PA-FABP), striated muscle and heart (H-FABP, mammary-derived growth inhibitor or MDGI), intestine (I-FABP), liver (L-FABP), myelin (M-FABP), and testis (T-FABP). Epithelial fatty acid-binding protein (E-FABP) binds stearic acid and may play a role in keratinocyte differentiation. E-FABP is upregulated in rat dorsal root ganglia after sciatic nerve crush and in differentiating neurons during development.

REFERENCES

- 1. Veerkamp, J.H. and Maatman, R.G. 1995. Cytoplasmic fatty acid-binding proteins: their structure and genes. Prog. Lipid Res. 34: 17-52.
- 2. Hotamisligil, G.S., et al. 1996. Uncoupling of obesity from Insulin resistance through a targeted mutation in aP2, the adipocyte fatty acid binding protein. Science 274: 1377-1379.
- 3. Storch, J. and Thumser, A.E. 2000. The fatty acid transport function of fatty acid-binding proteins. Biochim. Biophys. Acta 1486: 28-44.
- 4. Allen, G.W., et al. 2000. Depletion of a fatty acid-binding protein impairs neurite outgrowth in PC-12 cells. Brain Res. Mol. Brain Res. 76: 315-324.
- 5. Glatz, J.F. and Storch, J. 2001. Unravelling the significance of cellular fatty acid-binding proteins. Curr. Opin. Lipidol. 12: 267-274.
- 6. Veerkamp, J.H. and Zimmerman, A.W. 2001. Fatty acid-binding proteins of nervous tissue, J. Mol. Neurosci, 16: 133-142.

CHROMOSOMAL LOCATION

Genetic locus: FABP5 (human) mapping to 8q21.13.

SOURCE

E-FABP (D-12) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 75-109 within an internal region of E-FABP of human origin.

PRODUCT

Each vial contains 200 μ g lgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-373788 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

E-FABP (D-12) is recommended for detection of E-FABP of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

E-FABP (D-12) is also recommended for detection of E-FABP in additional species, including equine, canine and bovine.

Suitable for use as control antibody for E-FABP siRNA (h): sc-41237, E-FABP shRNA Plasmid (h): sc-41237-SH and E-FABP shRNA (h) Lentiviral Particles: sc-41237-V.

Molecular Weight of E-FABP: 15 kDa.

Positive Controls: E-FABP (h): 293T Lysate: sc-113029 or HEL 92.1.7 cell lysate: sc-2270.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG K BP-HRP: sc-516102 or m-IgG K BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). Immunofluorescence: use m-lgGκ BP-FITC: sc-516140 or m-lgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-lgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA







E-FABP (D-12): sc-373788. Immunoperoxidase staining of formalin fixed, paraffin-embedded human cervix uterine tissue showing cytoplasmic staining of squamous epithelial cells. Blocked with 0.25X UltraCruz® Blocking Reagent: sc-516214. Detected with m-lgG_K BP-B: sc-516142 and ImmunoCruz[®] ABC Kit: sc-516216.

RESEARCH USE

For research use only, not for use in diagnostic procedures.