

DNA pol α (D-4): sc-373884

BACKGROUND

DNA replication, recombination and repair, all of which are necessary for genomic stability, require the presence of exonucleases. In DNA replication, these enzymes are involved in the processing of Okazaki fragments, whereas in DNA repair, they function to excise damaged DNA fragments and correct recombinational mismatches. These exonucleases include the family of DNA polymerases. DNA pol α , β , δ , and ϵ are involved in DNA replication and repair. DNA pol δ and DNA pol ϵ are multisubunit enzymes, with DNA pol δ consisting of two subunits: p125, which interacts with the sliding DNA clamp protein PCNA; and p50. The nuclear-encoded DNA pol γ is the only DNA polymerase required for the replication of the mitochondrial DNA. DNA pol ζ is ubiquitously expressed in various tissues and mediates the cellular mechanism of damage-induced mutagenesis. DNA pol θ is a DNA polymerase-helicase that binds ATP and is involved in the repair of interstrand crosslinks.

CHROMOSOMAL LOCATION

Genetic locus: POLA1 (human) mapping to Xp22.11; Pola1 (mouse) mapping to X C3.

SOURCE

DNA pol α (D-4) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 3-31 near the N-terminus of DNA pol α of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-373884 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

DNA pol α (D-4) is recommended for detection of DNA pol α of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for DNA pol α siRNA (h): sc-37771, DNA pol α siRNA (m): sc-37772, DNA pol α shRNA Plasmid (h): sc-37771-SH, DNA pol α shRNA Plasmid (m): sc-37772-SH, DNA pol α shRNA (h) Lentiviral Particles: sc-37771-V and DNA pol α shRNA (m) Lentiviral Particles: sc-37772-V.


Molecular Weight of DNA pol α : 180 kDa.

Positive Controls: HeLa nuclear extract: sc-2120, Jurkat nuclear extract: sc-2132 or K-562 nuclear extract: sc-2130.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



DNA pol α (D-4): sc-373884. Western blot analysis of DNA pol α expression in HeLa (A), Jurkat (B) and K-562 (C) nuclear extracts.



DNA pol α (D-4): sc-373884. Immunofluorescence staining of methanol-fixed HeLa cells showing nuclear localization.

SELECT PRODUCT CITATIONS

- Tang, L., et al. 2019. DNA polymerase α is essential for intracellular amplification of hepatitis B virus covalently closed circular DNA. *PLoS Pathog.* 15: e1007742.
- Prokic, I, et al. 2020. Differential physiological role of BIN1 isoforms in skeletal muscle development, function and regeneration. *Dis. Model. Mech.* 13: dmm044354.
- Tsao, W.C., et al. 2021. Overexpression of oncogenic H-Ras in hTERT-immortalized and SV40-transformed human cells targets replicative and specialized DNA polymerases for depletion. *PLoS ONE* 16: e0251188.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.