

# DAP-3 (F-12): sc-373911

## BACKGROUND

The death-associated protein 3 (DAP-3), is a nucleotide-binding protein that contains a potential P-loop motif. It is a positive mediator of programmed cell death; overexpressed intact full-length protein is required in order to induce apoptosis. DAP-3 functions downstream of its receptor signaling complex and its death promoting effects depend on caspase activity. It also interacts with the glucocorticoid receptor (GR), where its main interaction domain is the amino-terminal region, which acts in a dominant-negative fashion to protect cells from apoptosis. DAP-3 protein may also play a role in modulating the cytoplasmic GR/HSP 90 complex. It is conserved at the functional, as well as the structural level, and is ubiquitously expressed in highly proliferative epithelial compartments of various tissues. Unlike a number of other proteins, DAP-3 retains its mitochondrial localization during the induction of apoptosis.

## REFERENCES

1. Kissil, J.L., et al. 1995. Isolation of DAP-3, a novel mediator of interferon- $\gamma$ -induced cell death. *J. Biol. Chem.* 270: 27932-27936.
2. Levy-Strumpf, N., et al. 1998. Death associated proteins (DAPs): from gene identification to the analysis of their apoptotic and tumor suppressive functions. *Oncogene* 17: 3331-3340.
3. Kissil, J.L., et al. 1999. Structure-function analysis of an evolutionary conserved protein, DAP-3, which mediates TNF $\alpha$ - and FAS-induced cell death. *EMBO J.* 18: 353-362.

## CHROMOSOMAL LOCATION

Genetic locus: DAP3 (human) mapping to 1q22; Dap3 (mouse) mapping to 3 F1.

## SOURCE

DAP-3 (F-12) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 63-95 within an internal region of DAP-3 of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG $_1$  kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

DAP-3 (F-12) is available conjugated to agarose (sc-373911 AC), 500  $\mu$ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-373911 HRP), 200  $\mu$ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-373911 PE), fluorescein (sc-373911 FITC), Alexa Fluor<sup>®</sup> 488 (sc-373911 AF488), Alexa Fluor<sup>®</sup> 546 (sc-373911 AF546), Alexa Fluor<sup>®</sup> 594 (sc-373911 AF594) or Alexa Fluor<sup>®</sup> 647 (sc-373911 AF647), 200  $\mu$ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor<sup>®</sup> 680 (sc-373911 AF680) or Alexa Fluor<sup>®</sup> 790 (sc-373911 AF790), 200  $\mu$ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-373911 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

Alexa Fluor<sup>®</sup> is a trademark of Molecular Probes, Inc., Oregon, USA

## APPLICATIONS

DAP-3 (F-12) is recommended for detection of DAP-3 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for DAP-3 siRNA (h): sc-37381, DAP-3 siRNA (m): sc-155880, DAP-3 shRNA Plasmid (h): sc-37381-SH, DAP-3 shRNA Plasmid (m): sc-155880-SH, DAP-3 shRNA (h) Lentiviral Particles: sc-37381-V and DAP-3 shRNA (m) Lentiviral Particles: sc-155880-V.


Molecular Weight of DAP-3: 46 kDa.

Positive Controls: NCI-H929 whole cell lysate: sc-364786, HeLa whole cell lysate: sc-2200 or MCF7 whole cell lysate: sc-2206.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

## DATA



DAP-3 (F-12): sc-373911. Western blot analysis of DAP-3 expression in HeLa (A), NCI-H929 (B) and MCF7 (C) whole cell lysates.



DAP-3 (F-12): sc-373911. Immunofluorescence staining of methanol-fixed HeLa cells showing nuclear and cytoplasmic localization.

## SELECT PRODUCT CITATIONS

1. Sui, L., et al. 2023. Death associated protein-3 (DAP3) and DAP3 binding cell death enhancer-1 (DELE1) in human colorectal cancer, and their impacts on clinical outcome and chemoresistance. *Int. J. Oncol.* 62: 7.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.