

## CTGF (B-6): sc-373936



The Power to Question

## BACKGROUND

Connective tissue growth factor (CTGF, also known as hypertrophic chondrocyte-specific gene product 24 or Hcs24), is a member of the CCN family of immediate early proteins, which are involved in cell proliferation, migration and matrix production. CTGF is a cysteine-rich peptide that is secreted by endothelial cells, fibroblasts, smooth muscle cells and myofibroblasts. Its expression is increased in various human and animal fibrotic diseases. Specifically, CTGF has been observed to be strongly upregulated in human proliferative and fibrogenic renal disease. In addition, CTGF is a growth factor for vascular smooth muscle cells (VSMC) and may play a similar role in promoting VSMC growth and migration *in vitro*.

## CHROMOSOMAL LOCATION

Genetic locus: CTGF (human) mapping to 6q23.2; Ctgf (mouse) mapping to 10 A4.

## SOURCE

CTGF (B-6) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 170-207 within an internal region of CTGF of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>3</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

CTGF (B-6) is available conjugated to agarose (sc-373936 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-373936 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-373936 PE), fluorescein (sc-373936 FITC), Alexa Fluor® 488 (sc-373936 AF488), Alexa Fluor® 546 (sc-373936 AF546), Alexa Fluor® 594 (sc-373936 AF594) or Alexa Fluor® 647 (sc-373936 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-373936 AF680) or Alexa Fluor® 790 (sc-373936 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-373936 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

## APPLICATIONS

CTGF (B-6) is recommended for detection of CTGF of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

CTGF (B-6) is also recommended for detection of CTGF in additional species, including equine, canine and bovine.


Suitable for use as control antibody for CTGF siRNA (h): sc-39329, CTGF siRNA (m): sc-39330, CTGF shRNA Plasmid (h): sc-39329-SH, CTGF shRNA Plasmid (m): sc-39330-SH, CTGF shRNA (h) Lentiviral Particles: sc-39329-V and CTGF shRNA (m) Lentiviral Particles: sc-39330-V.

Molecular Weight of CTGF: 38 kDa.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## DATA



CTGF (B-6): sc-373936. Western blot analysis of CTGF expression in non-transfected: sc-117752 (A) and human CTGF transfected: sc-111612 (B) 293T whole cell lysates.

## SELECT PRODUCT CITATIONS

- Ma, L., et al. 2014. Cluster of differentiation 166 (CD166) regulated by phosphatidylinositol 3-kinase (PI3K)/Akt signaling to exert its anti-apoptotic role via yes-associated protein (YAP) in liver cancer. *J. Biol. Chem.* 289: 6921-6933.
- Tao, Y., et al. 2018. Huaier augmented the chemotherapeutic sensitivity of oxaliplatin via downregulation of YAP in hepatocellular carcinoma. *J. Cancer* 9: 3962-3970.
- Wu, M., et al. 2019. Reduced asymmetric dimethylarginine accumulation through inhibition of the type I protein arginine methyltransferases promotes renal fibrosis in obstructed kidneys. *FASEB J.* 33: 6948-6956.
- Xu, L., et al. 2020. Emodin ameliorates tubulointerstitial fibrosis in obstructed kidneys by inhibiting EZH2. *Biochem. Biophys. Res. Commun.* 534: 279-285.
- Bansod, S., et al. 2021. Inhibition of discoidin domain receptors by imatinib prevented pancreatic fibrosis demonstrated in experimental chronic pancreatitis model. *Sci. Rep.* 11: 12894.
- Namba, M., et al. 2021. Creation of X-linked Alport syndrome rat model with Col4a5 deficiency. *Sci. Rep.* 11: 20836.
- Yang, L., et al. 2022. Elucidating the novel mechanism of ligustrazine in preventing postoperative peritoneal adhesion formation. *Oxid. Med. Cell. Longev.* 2022: 9226022.
- Marroncini, G., et al. 2022. Hyponatremia-related liver steatofibrosis and impaired spermatogenesis: evidence from a mouse model of the syndrome of inappropriate antidiuresis. *J. Endocrinol. Invest.* 46: 967-983.
- Qian, Z., et al. 2023. Bufalin inhibits the proliferation of lung cancer cells by suppressing Hippo-YAP pathway. *Cell. Signal.* 109: 110746.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA