# SANTA CRUZ BIOTECHNOLOGY, INC.

# IL-2Rα (H-6): sc-374034



### BACKGROUND

The IL-2 receptor is a multicomponent complex consisting of three subunits,  $\alpha$ ,  $\beta$  and  $\gamma$ , each of which is required for high-affinity binding of IL-2. The  $\alpha$  chain functions primarily in binding IL-2, whereas the  $\beta$  and  $\gamma$  chains contribute to IL-2 binding and are essential to IL-2-induced activation of signaling pathways leading to T cell growth. Both IL-4R and IL-7R were initially described as single chain, high-affinity, ligand-binding cytokine receptors. However, it is now well-established that the IL-2R $\gamma$  chain functions as a second subunit of the high-affinity IL-4R and IL-7R receptors. Consequently, the originally described subunits of these latter receptors are now referred to as IL-4R $\alpha$  and IL-7R $\alpha$ , respectively, while the common subunit is referred to as  $\gamma$ c. Although the common  $\gamma$  chain enhances ligand binding in these three cytokine receptors, it has no capacity to bind these ligands on its own. There is evidence that the  $\gamma$ c chain is also a subunit of IL-13R.

### REFERENCES

- 1. Paterson, D.J., et al. 1987. Antigens of activated rat T lymphocytes including a molecule of 50,000  $\rm M_r$  detected only on CD4+ T blasts. Mol. Immunol. 24: 1281-1290.
- Mosley, B., et al. 1989. The murine interleukin-4 receptor: molecular cloning and characterization of secreted and membrane-bound forms. Cell 59: 335-348.
- Goodwin, R.G., et al. 1990. Cloning of the human and murine interleukin-7 receptors: demonstration of a soluble form and homology to a new receptor superfamily. Cell 60: 941-951.
- 4. Takeshita, T., et al. 1992. Cloning of the  $\gamma$  chain of the human IL-2 receptor. Science 257: 379-382.
- 5. Cao, X., et al. 1993. Characterization of cDNAs encoding the murine interleukin-2 receptor (IL-2R)  $\gamma$  chain: chromosomal mapping and tissue specificity of IL-2R  $\gamma$  chain expression. Proc. Natl. Acad. Sci. USA 90: 8464-8468.
- 6. Kondo, M., et al. 1993. Sharing of the interleukin-2 (IL-2) receptor  $\gamma$  chain between receptors for IL-2 and IL-4. Science 262: 1874-1877.
- 7. Russell, S.M., et al. 1993. Interleukin-2 receptor  $\gamma$  chain: a functional component of the interleukin-4 receptor. Science 262: 1880-1883.
- 8. Minami, Y., et al. 1993. The IL-2 receptor complex: its structure, function and target genes. Annu. Rev. Immunol. 11: 245-268.

#### **CHROMOSOMAL LOCATION**

Genetic locus: IL2RA (human) mapping to 10p15.1.

#### SOURCE

IL-2R $\alpha$  (H-6) is a mouse monoclonal antibody raised against amino acids 1-272 representing full length IL-2R $\alpha$  of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG<sub>2a</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

### **APPLICATIONS**

IL-2R $\alpha$  (H-6) is recommended for detection of IL-2R $\alpha$  of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for IL-2R $\alpha$  siRNA (h): sc-29367, IL-2R $\alpha$  shRNA Plasmid (h): sc-29367-SH and IL-2R $\alpha$  shRNA (h) Lentiviral Particles: sc-29367-V.

Molecular Weight of IL-2R $\alpha$ : 55 kDa.

Positive Controls: HuT 78 whole cell lysate: sc-2208 or IL-2R  $\alpha$  (h2): 293T Lysate: sc-176584.

## **RECOMMENDED SUPPORT REAGENTS**

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG K BP-HRP: sc-516102 or m-IgG K BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG K BP-FITC: sc-516140 or m-IgG K BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

#### DATA



#### **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

#### PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.