

CRM1 (A-7): sc-374124

BACKGROUND

Protein transport across the nucleus is a selective, multistep process involving several cytoplasmic factors. Proteins must be recognized as import substrates, dock at the nuclear pore complex and translocate across the nuclear envelope in an ATP-dependent fashion. Two cytosolic factors centrally involved in the recognition and docking process are the karyopherin α 1 and karyopherin β 1 subunits. p62 glycoprotein is a nucleoporin that is not only involved in the nuclear import of proteins, but also the export of nascent mRNA strands. NTF2 (nuclear transport factor 2) interacts with nucleoporin p62 as a homodimer composed of two monomers, and may be an obligate component of functional p62. CRM1 has been shown to be an export receptor for leucine-rich proteins that contain the nuclear export signal (NES).

CHROMOSOMAL LOCATION

Genetic locus: XPO1 (human) mapping to 2p15; Xpo1 (mouse) mapping to 11 A3.2.

SOURCE

CRM1 (A-7) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 1043-1071 at the C-terminus of CRM1 of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-374124 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

CRM1 (A-7) is recommended for detection of CRM1 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CRM1 siRNA (h): sc-35116, CRM1 siRNA (m): sc-35117, CRM1 shRNA Plasmid (h): sc-35116-SH, CRM1 shRNA Plasmid (m): sc-35117-SH, CRM1 shRNA (h) Lentiviral Particles: sc-35116-V and CRM1 shRNA (m) Lentiviral Particles: sc-35117-V.


Molecular Weight of CRM1: 115 kDa.

Positive Controls: HeLa nuclear extract: sc-2120, A-431 nuclear extract: sc-2122 or K-562 nuclear extract: sc-2130.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



CRM1 (A-7): sc-374124. Western blot analysis of CRM1 expression in HeLa (A) and K-562 (B) nuclear extracts.



CRM1 (A-7): sc-374124. Immunofluorescence staining of methanol-fixed HeLa cells showing nuclear localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human colon tissue showing nuclear and cytoplasmic staining of glandular cells and nuclear staining of endothelial cells (B).

SELECT PRODUCT CITATIONS

1. Martin, A.P., et al. 2019. STK38 kinase acts as XPO1 gatekeeper regulating the nuclear export of autophagy proteins and other cargoes. EMBO Rep. 20: e48150.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **CRM1 (C-1): sc-74454** for CRM1 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.