

LDH-D (G-8): sc-374127

BACKGROUND

The lactate dehydrogenase family (LDH) consists of three members, designated LDH-A, LDH-B and LDH-C, all of which work in concert to catalyze the final step of anaerobic glycolysis, namely the conversion of L-lactate and NAD⁺ to pyruvate and NADH. Each family member displays a specific tissue distribution pattern, with LDH-A present in muscle and LDH-B present in heart, while LDH-C expression is confined to testes and sperm. A fourth possible member, termed LDH-D (lactate dehydrogenase D), probable D-lactate dehydrogenaseor DLD, is a 507 amino acid mitochondrial protein belonging to the D-isomer specific 2-hydroxyacid dehydrogenase family. Existing as two alternatively spliced isoforms, LDH-D is moderately expressed in liver and heart with lower levels found in kidney and skeletal muscle.

REFERENCES

1. Edwards, Y.H., Povey, S., LeVan, K.M., Driscoll, C.E., Millan, J.L. and Goldberg, E. 1987. Locus determining the human sperm-specific lactate dehydrogenase, LDHC, is syntenic with LDHA. *Dev. Genet.* 8: 219-232.
2. LeVan, K.M. and Goldberg, E. 1991. Properties of human testis-specific lactate dehydrogenase expressed from *Escherichia coli*. *Biochem. J.* 273: 587-592.
3. Kanno, T. and Maekawa, M. 1995. Lactate dehydrogenase M-subunit deficiencies: clinical features, metabolic background, and genetic heterogeneities. *Muscle Nerve Suppl.* 3: S54-S60.
4. Kopperschläger, G. and Kirchberger, J. 1996. Methods for the separation of lactate dehydrogenases and clinical significance of the enzyme. *J. Chromatogr. B, Biomed. Appl.* 684: 25-49.
5. Niwakawa, M. and Tobisu, K. 2001. The role of tumor markers in the treatment of germ cell tumor. *Gan To Kagaku Ryoho* 28: 1159-1165.
6. Flick, M.J. and Konieczny, S.F. 2002. Identification of putative mammalian D-lactate dehydrogenase enzymes. *Biochem. Biophys. Res. Commun.* 295: 910-916.
7. Online Mendelian Inheritance in Man, OMIM™. 2009. Johns Hopkins University, Baltimore, MD. MIM Number: 607490. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: LDHD (human) mapping to 16q23.1; Ldhd (mouse) mapping to 8 E1.

SOURCE

LDH-D (G-8) is a mouse monoclonal antibody raised against amino acids 1-195 mapping at the N-terminus of LDH-D of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

LDH-D (G-8) is recommended for detection of LDH-D of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for LDH-D siRNA (h): sc-93282, LDH-D siRNA (m): sc-146691, LDH-D shRNA Plasmid (h): sc-93282-SH, LDH-D shRNA Plasmid (m): sc-146691-SH, LDH-D shRNA (h) Lentiviral Particles: sc-93282-V and LDH-D shRNA (m) Lentiviral Particles: sc-146691-V.


Molecular Weight of LDH-D: 55 kDa.

Positive Controls: mouse heart extract: sc-2254, human heart extract: sc-363763 or LADMAC whole cell lysate: sc-364189.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



LDH-D (G-8): sc-374127. Western blot analysis of LDH-D expression in human heart (A), mouse heart (B), human skeletal muscle (C) and mouse skeletal muscle (D) tissue extracts and LADMAC whole cell lysate (E).



LDH-D (G-8): sc-374127. Immunoperoxidase staining of formalin fixed, paraffin-embedded human heart muscle tissue showing cytoplasmic staining of myocytes.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.