

eIF3 β (A-8): sc-374155

BACKGROUND

The initiation of protein synthesis in eukaryotic cells is regulated by interactions between protein initiation factors and RNA molecules. Eukaryotic initiation factors (eIFs) are utilized in a sequence of reactions that lead to 80S ribosomal assembly and, ultimately, translation. The eukaryotic initiation factor-3 (eIF3) scaffolding structure is the largest of the eIF complexes and includes eIF3 α , eIF3 β , eIF3 γ , eIF3 δ , eIF3 ϵ , eIF3 ζ , eIF3 η and eIF3 θ , all of which function to control the assembly of the 40S ribosomal subunit. Association of eIF3 proteins with the 40S ribosomal subunit stabilizes eIF2-GTP-Met-tRNA^{iMet} complex association and mRNA binding, and promotes dissociation of 80S ribosomes into 40S and 60S subunits, thereby promoting the assembly of the pre-initiation complex. Overexpression of eIF3 proteins is common in several cancers, suggesting a role for eIF3 proteins in tumorigenesis.

REFERENCES

- Valásek, L., et al. 2004. Interactions of eukaryotic translation initiation factor 3 (eIF3) subunit NIP1/c with eIF1 and eIF5 promote preinitiation complex assembly and regulate start codon selection. *Mol. Cell. Biol.* 24: 9437-9455.
- Peterson, T.R., et al. 2005. eIF3: a connectOR of S6K1 to the translation preinitiation complex. *Mol. Cell* 20: 655-657.
- Dong, Z., et al. 2006. Initiation factor eIF3 and regulation of mRNA translation, cell growth, and cancer. *Crit. Rev. Oncol. Hematol.* 59: 169-180.
- LeFebvre, A.K., et al. 2006. Translation initiation factor eIF4G-1 binds to eIF3 through the eIF3 ϵ subunit. *J. Biol. Chem.* 281: 22917-22932.
- Hinnebusch, A.G. 2006. eIF3: a versatile scaffold for translation initiation complexes. *Trends Biochem. Sci.* 31: 553-562.
- Masutani, M., et al. 2007. Reconstitution reveals the functional core of mammalian eIF3. *EMBO J.* 26: 3373-3383.
- Zhang, L., et al. 2007. Individual overexpression of five subunits of human translation initiation factor eIF3 promotes malignant transformation of immortal fibroblast cells. *J. Biol. Chem.* 282: 5790-5800.
- Sato, H., et al. 2007. Measles virus N protein inhibits host translation by binding to eIF3-p40. *J. Virol.* 81: 11569-11576.
- Zhang, L., et al. 2008. An oncogenic role for the phosphorylated η -subunit of human translation initiation factor eIF3. *J. Biol. Chem.* 283: 24047-24060.

CHROMOSOMAL LOCATION

Genetic locus: EIF3I (human) mapping to 1p35.1; Eif3i (mouse) mapping to 4 D2.2.

SOURCE

eIF3 β (A-8) is a mouse monoclonal antibody raised against amino acids 1-300 mapping within an internal region of eIF3 β of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

eIF3 β (A-8) is recommended for detection of eIF3 β of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for eIF3 β siRNA (h): sc-60080, eIF3 β siRNA (m): sc-60081, eIF3 β shRNA Plasmid (h): sc-60080-SH, eIF3 β shRNA Plasmid (m): sc-60081-SH, eIF3 β shRNA (h) Lentiviral Particles: sc-60080-V and eIF3 β shRNA (m) Lentiviral Particles: sc-60081-V.


Molecular Weight of eIF3 β : 36 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, Hep G2 cell lysate: sc-2227 or Jurkat whole cell lysate: sc-2204.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA



eIF3 β (A-8): sc-374155. Western blot analysis of eIF3 β expression in HeLa (A), Jurkat (B), Hep G2 (C), NIH/3T3 (D), AMJ2-C8 (E) and Neuro-2A (F) whole cell lysates.



eIF3 β (A-8): sc-374155. Western blot analysis of eIF3 β expression in T24 (A), SJRH30 (B), Jurkat (C) and MCF7 (D) whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.