

Geminin (H-3): sc-374187

BACKGROUND

Geminin is a nuclear protein that regulates the initiation of DNA replication during the cell cycle. DNA replication requires the coordinated association of cdc6 and minichromosome maintenance (MCM) proteins with chromatin. Geminin blocks this assembly of the MCM into the prereplication complex and, in turn, prevents replication from occurring. Expression of Geminin fluctuates throughout the cell cycle with Geminin levels lowest at G₁. Throughout S, G₂ and M phases, Geminin levels are consistently elevated followed by a decrease during mitosis. The initiation of DNA replication is dependent on the degradation of Geminin during mitosis and the absence of Geminin throughout G₁ phase. Geminin degradation is mediated by the anaphase-promoting complex (APC), which specifically targets B-type cyclins and other proteins containing a destruction box motif for degradation by ubiquitin-mediated proteolysis.

REFERENCES

1. Yu, H., et al. 1996. Identification of a novel ubiquitin-conjugating enzyme involved in mitotic cyclin degradation. *Curr. Biol.* 6: 455-466.
2. Rowles, A., et al. 1997. Chromatin proteins involved in the initiation of DNA replication. *Curr. Opin. Genet. Dev.* 7: 152-157.
3. Liang, C., et al. 1997. Persistent initiation of DNA replication and chromatin-bound MCM proteins during the cell cycle in cdc6 mutants. *Genes Dev.* 11: 3375-3386.
4. Page, A.M., et al. 1997. The anaphase promoting complex. *Cancer Surv.* 29: 133-150.
5. McGarry, T.J., et al. 1998. Geminin, an inhibitor of DNA replication, is degraded during mitosis. *Cell* 93: 1043-1053.
6. Kroll, K.L., et al. 1998. Geminin, a neuralizing molecule that demarcates the future neural plate at the onset of gastrulation. *Development* 125: 3247-3258.

CHROMOSOMAL LOCATION

Genetic locus: Gmn (mouse) mapping to 13 A3.1.

SOURCE

Geminin (H-3) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 2-29 at the N-terminus of Geminin of mouse origin.

PRODUCT

Each vial contains 200 µg IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-374187 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Geminin (H-3) is recommended for detection of Geminin of mouse and rat origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Geminin siRNA (m): sc-108025, Geminin shRNA Plasmid (m): sc-108025-SH and Geminin shRNA (m) Lentiviral Particles: sc-108025-V.


Molecular Weight of Geminin: 35 kDa.

Positive Controls: MM-142 cell lysate: sc-2246, Geminin (m): 293T Lysate: sc-120468 or MM-142 nuclear extract: sc-2139.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



Geminin (H-3): sc-374187. Western blot analysis of Geminin expression in non-transfected: sc-117752 (A) and mouse Geminin transfected: sc-120468 (B) 293T whole cell lysates.



Geminin (H-3): sc-374187. Immunofluorescence staining of methanol-fixed NIH/3T3 cells showing nuclear localization.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.