

CCS (D-7): sc-374205

BACKGROUND

Cu-Zn superoxide dismutase-1 (SOD-1) is a well characterized cytosolic scavenger of oxygen free radicals that requires copper and zinc binding to potentiate its enzymatic activity. Copper chaperone for SOD-1 (CCS) is essential for the incorporation of copper into SOD-1, and therefore is necessary for its enzymatic activity. CCS prevents copper ions from binding to intracellular copper scavengers and provides the SOD-1 enzyme with the necessary copper cofactor. CCS escorts copper only to SOD-1 and fails to deliver copper to proteins in the mitochondria, nucleus or secretory pathway. CCS interacts with both wildtype and mutated forms of SOD-1 through CCS domains that are homologous in SOD-1. CCS exists as a homodimer that may form a heterodimer with SOD-1 during copper loading. While many tissues express CCS, the chaperone is most abundant in the kidney, liver and Purkinje cells in the neurophil of the central nervous system.

CHROMOSOMAL LOCATION

Genetic locus: CCS (human) mapping to 11q13.2; Ccs (mouse) mapping to 19 A.

SOURCE

CCS (D-7) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 199-235 within an internal region of CCS of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-374205 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

CCS (D-7) is recommended for detection of CCS of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

CCS (D-7) is also recommended for detection of CCS in additional species, including porcine.

Suitable for use as control antibody for CCS siRNA (h): sc-29956, CCS siRNA (m): sc-29957, CCS shRNA Plasmid (h): sc-29956-SH, CCS shRNA Plasmid (m): sc-29957-SH, CCS shRNA (h) Lentiviral Particles: sc-29956-V and CCS shRNA (m) Lentiviral Particles: sc-29957-V.


Molecular Weight of CCS: 35 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, HL-60 whole cell lysate: sc-2209 or Hep G2 cell lysate: sc-2227.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



CCS (D-7): sc-374205. Western blot analysis of CCS expression in HL-60 (A), Hep G2 (B), HeLa (C) and c4 (D) whole cell lysates.



CCS (D-7): sc-374205. Immunoperoxidase staining of formalin fixed, paraffin-embedded human tonsil tissue showing cytoplasmic and nuclear staining of cells in non-germinal center.

SELECT PRODUCT CITATIONS

- Ghaffari, R., et al. 2019. Copper transporter 1 (CTR1) expression by mouse testicular germ cells, but not Sertoli cells, is essential for functional spermatogenesis. PLoS ONE 14: e0215522.
- Kim, J.H., et al. 2021. Lysosomal SLC46A3 modulates hepatic cytosolic copper homeostasis. Nat. Commun. 12: 290.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.