SANTA CRUZ BIOTECHNOLOGY, INC.

IMPA1 (H-7): sc-374234



BACKGROUND

IMPA1, also known as Myo-inositol monophosphatase 1, is responsible for the procurement of inositol that is required for synthesis of phosphatidylinositol and polyphosphoinositides. IMPA1 exists as a homodimer and has been identified as the pharmacological target for lithium action in the brain. IMPA1 is the principal enzyme of the phosphatidyl inositol signaling pathway, and inhibition of inositol monophosphatase hydrolysis may underlie the anti-manic and anti-depressant actions of Li⁺. Studies indicate that a variation in the 277 codon coding region of the IMPA1 gene has not been observed in manic-depressive patients, therefore suggesting that polymorphisms or mutations in the noncoding regions of this gene may influence the lithium response in psychiatric patients.

REFERENCES

- McAllister, G., et al. 1992. cDNA cloning of human and rat brain Myoinositol monophosphatase. Expression and characterization of the human recombinant enzyme. Biochem. J. 284: 749-754.
- Klein, P.S., et al. 1996. A molecular mechanism for the effect of lithium on development. Proc. Natl. Acad. Sci. USA 93: 8455-8459.
- Steen, V.M., et al. 1996. Lack of genetic variation in the coding region of the Myo-inositol monophosphatase gene in lithium-treated patients with manic depressive illness. Pharmacogenetics 6: 113-116.
- Sjoholt, G., et al. 1997. Genomic structure and chromosomal localization of a human Myo-inositol monophosphatase gene (IMPA). Genomics 45: 113-122.

CHROMOSOMAL LOCATION

Genetic locus: IMPA1 (human) mapping to 8q21.13; Impa1 (mouse) mapping to 3 A1.

SOURCE

IMPA1 (H-7) is a mouse monoclonal antibody raised against amino acids 1-85 mapping at the N-terminus of IMPA1 of human origin.

PRODUCT

Each vial contains 200 $\mu g\, lgG_1$ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

IMPA1 (H-7) is available conjugated to agarose (sc-374234 AC), 500 µg/ 0.25 ml agarose in 1 ml, for IP; to HRP (sc-374234 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-374234 PE), fluorescein (sc-374234 FITC), Alexa Fluor[®] 488 (sc-374234 AF488), Alexa Fluor[®] 546 (sc-374234 AF546), Alexa Fluor[®] 594 (sc-374234 AF594) or Alexa Fluor[®] 647 (sc-374234 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-374234 AF680) or Alexa Fluor[®] 790 (sc-374234 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor $^{\circ}$ is a trademark of Molecular Probes, Inc., Oregon, USA

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

IMPA1 (H-7) is recommended for detection of IMPA1 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), isotochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for IMPA1 siRNA (h): sc-61115, IMPA1 siRNA (m): sc-61116, IMPA1 shRNA Plasmid (h): sc-61115-SH, IMPA1 shRNA Plasmid (m): sc-61116-SH, IMPA1 shRNA (h) Lentiviral Particles: sc-61115-V and IMPA1 shRNA (m) Lentiviral Particles: sc-61116-V.

Molecular Weight of IMPA1: 29 kDa.

Positive Controls: IMR-32 cell lysate: sc-2409, C6 whole cell lysate: sc-364373 or Neuro-2A whole cell lysate: sc-364185.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA





IMPA1 (H-7): sc-374234. Western blot analysis of IMPA1 expression in IMR-32 (\mathbf{A}), Neuro-2A (\mathbf{B}) and C6 (\mathbf{C}) whole cell lysates and human cerebral cortex tissue extract (\mathbf{D}).

IMPA1 (H-7): sc-374234. Immunoperoxidase staining of formalin fixed, paraffin-embedded human thyroid tissue showing nuclear and cytoplasmic staining of glandular cells.

SELECT PRODUCT CITATIONS

 Hsu, C.C., et al. 2021. Inositol serves as a natural inhibitor of mitochondrial fission by directly targeting AMPK. Mol. Cell 81: 3803-3819.e7.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.