

ZAP-70 (A-8): sc-374347

BACKGROUND

The activation of T lymphocytes by antigens is mediated by the T cell receptor (TCR) which is a multisubunit complex assembled from at least six different genes. The TCR subunits include the T α and β chains, the CD3 γ , δ and ϵ chains and a ζ -containing homodimer or heterodimer. The disulfide-linked T α - β heterodimer is responsible for antigen recognition, but the short five amino acid cytoplasmic domains of T α and β are unlikely to be sufficient to couple to intracellular signaling pathways. In contrast, the structured features of the CD3 and ζ subunits suggest a role in signal transduction. Of these, the ζ chain, which is expressed as either a homodimer or heterodimer, has a short extracellular domain of only nine amino acids, but a larger 113 amino acid cytoplasmic domain. A tyrosine phosphoprotein, ZAP-70, has been identified that associates with ζ and undergoes tyrosine phosphorylation following TCR stimulation.

REFERENCES

1. Clevers, H., et al. 1988. The T cell receptor/CD3 complex: a dynamic protein ensemble. *Annu. Rev. Immunol.* 6: 629-662.
2. Baniyash, M., et al. 1988. Disulfide linkage of the ζ and η chains of the T cell receptor. Possible identification of two structural classes of receptors. *J. Biol. Chem.* 263: 9874-9878.
3. Baniyash, M., et al. 1988. The T cell antigen receptor ζ chain is tyrosine phosphorylated upon activation. *J. Biol. Chem.* 263: 18225-18230.
4. Baniyash, M., et al. 1989. The isolation and characterization of the murine T cell antigen receptor ζ chain gene. *J. Biol. Chem.* 264: 13252-13257.
5. Frank, S.J., et al. 1990. The structure and signaling function of the invariant T cell receptor components. *Semin. Immunol.* 2: 89-97.

CHROMOSOMAL LOCATION

Genetic locus: ZAP70 (human) mapping to 2q11.2; Zap70 (mouse) mapping to 1 B.

SOURCE

ZAP-70 (A-8) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 5-35 near the N-terminus of ZAP-70 of human origin.

PRODUCT

Each vial contains 200 μ g IgG γ_1 kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-374347 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

ZAP-70 (A-8) is recommended for detection of ZAP-70 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

ZAP-70 (A-8) is also recommended for detection of ZAP-70 in additional species, including bovine.

Suitable for use as control antibody for ZAP-70 siRNA (h): sc-29526, ZAP-70 siRNA (m): sc-36867, ZAP-70 shRNA Plasmid (h): sc-29526-SH, ZAP-70 shRNA Plasmid (m): sc-36867-SH, ZAP-70 shRNA (h) Lentiviral Particles: sc-29526-V and ZAP-70 shRNA (m) Lentiviral Particles: sc-36867-V.


Molecular Weight of ZAP-70: 70 kDa.

Positive Controls: HuT 78 whole cell lysate: sc-2208, Jurkat whole cell lysate: sc-2204 or MOLT-4 cell lysate: sc-2233.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



ZAP-70 (A-8): sc-374347. Western blot analysis of ZAP-70 expression in Jurkat whole cell lysate.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



See **ZAP-70 (1E7.2): sc-32760** for ZAP-70 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.