

SNM1B (F-8): sc-374351

BACKGROUND

DNA interstrand cross-links (ICLs) pose lethal threats to DNA as they inhibit segregation, replication and transcription. The mechanism of ICL repair is complex but is at least partly conserved between *Saccharomyces cerevisiae* and mammals. SNM1B (SNM1 homolog B), also known as DCLRE1B [DNA cross-link repair 1B (PSO2 homolog, *S. cerevisiae*)] or APOLLO, is a 532 amino acid nuclear protein that localizes to discrete foci and is likely required for DNA interstrand cross-link repair. SNM1B assists in the maintenance of telomeres during S-phase and interacts with TRF2 (telomeric repeat binding factor 2), a protein involved in telomeric organization and protection, in the early DNA-damage response. A member of the DNA repair metallo- β -lactamase (DRMBL) family, SNM1B becomes phosphorylated following translation, either by ATM or ATR, and is encoded by a gene located on human chromosome 1.

REFERENCES

1. Dronkert, M.L., et al. 2000. Disruption of mouse SNM1 causes increased sensitivity to the DNA interstrand cross-linking agent mitomycin C. *Mol. Cell. Biol.* 20: 4553-4561.
2. Demuth, I., et al. 2004. Human SNM1B is required for normal cellular response to both DNA interstrand crosslink-inducing agents and ionizing radiation. *Oncogene* 23: 8611-8618.
3. Freibaum, B.D. and Counter, C.M. 2006. hSNM1B is a novel telomere-associated protein. *J. Biol. Chem.* 281: 15033-15036.
4. Demuth, I., et al. 2008. Endogenous hSNM1B/Apollo interacts with TRF2 and stimulates ATM in response to ionizing radiation. *DNA Repair* 7: 1192-1201.

CHROMOSOMAL LOCATION

Genetic locus: DCLRE1B (human) mapping to 1p13.2.

SOURCE

SNM1B (F-8) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 419-451 within an internal region of SNM1B of human origin.

PRODUCT

Each vial contains 200 μ g IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

SNM1B (F-8) is available conjugated to agarose (sc-374351 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-374351 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-374351 PE), fluorescein (sc-374351 FITC), Alexa Fluor[®] 488 (sc-374351 AF488), Alexa Fluor[®] 546 (sc-374351 AF546), Alexa Fluor[®] 594 (sc-374351 AF594) or Alexa Fluor[®] 647 (sc-374351 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-374351 AF680) or Alexa Fluor[®] 790 (sc-374351 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-374351 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

SNM1B (F-8) is recommended for detection of SNM1B of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for SNM1B siRNA (h): sc-88509, SNM1B shRNA Plasmid (h): sc-88509-SH and SNM1B shRNA (h) Lentiviral Particles: sc-88509-V.


Molecular Weight of SNM1B: 60 kDa.

Positive Controls: SNM1B (h): 293T Lysate: sc-113949.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA



SNM1B (F-8): sc-374351. Western blot analysis of SNM1B expression in non-transfected: sc-117752 (A) and human SNM1B transfected: sc-113949 (B) 293T whole cell lysates.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA