

Ran BP-1 (D-8): sc-374352

BACKGROUND

The small Ras-related protein Ran, also called TC4, is a nuclear localized GTPase implicated in a diverse array of cellular processes including DNA replication, entry into and exit from mitosis and the transport of RNA and proteins through the nuclear pore complex. Like Ras, active Ran GTP and inactive Ran GDP levels are tightly regulated by guanine nucleotide exchange factors (GEFs) and GTPase activating proteins (GAPs). The abundant GEF, RCC1 (regulator of chromosome condensation 1), increases the rate at which Ran exchanges GDP for GTP. Ran GAP1 opposes the effects of RCC1 by increasing the rate at which Ran hydrolyzes GTP to GDP. A protein designated Ran BP1 has no intrinsic GAP activity, and functions as a GEF inhibitor deactivating RCC1 and thereby indirectly increasing the ratio of Ran GDP to Ran GTP. The Ran BP2 protein has been proposed as the Ran GTP docking site at the periphery of the nuclear pore complex.

CHROMOSOMAL LOCATION

Genetic locus: RANBP1 (human) mapping to 22q11.21; Ranbp1 (mouse) mapping to 16 A3.

SOURCE

Ran BP-1 (D-8) is a mouse monoclonal antibody raised against amino acids 159-203 mapping at the C-terminus of Ran BP-1 of mouse origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Ran BP-1 (D-8) is available conjugated to agarose (sc-374352 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-374352 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-374352 PE), fluorescein (sc-374352 FITC), Alexa Fluor® 488 (sc-374352 AF488), Alexa Fluor® 546 (sc-374352 AF546), Alexa Fluor® 594 (sc-374352 AF594) or Alexa Fluor® 647 (sc-374352 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-374352 AF680) or Alexa Fluor® 790 (sc-374352 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

APPLICATIONS

Ran BP-1 (D-8) is recommended for detection of Ran BP-1 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Ran BP-1 siRNA (h): sc-41848, Ran BP-1 siRNA (m): sc-41849, Ran BP-1 shRNA Plasmid (h): sc-41848-SH, Ran BP-1 shRNA Plasmid (m): sc-41849-SH, Ran BP-1 shRNA (h) Lentiviral Particles: sc-41848-V and Ran BP-1 shRNA (m) Lentiviral Particles: sc-41849-V.


Molecular Weight of Ran BP-1: 28 kDa.

Positive Controls: NIH/3T3 whole cell lysate: sc-2210, F9 cell lysate: sc-2245 or 3611-RF whole cell lysate: sc-2215.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



Ran BP-1 (D-8): sc-374352. Western blot analysis of Ran BP-1 expression in F9 (A), c4 (B), KNRK (C) and 3611-RF (D) whole cell lysates.



Ran BP-1 (D-8): sc-374352. Western blot analysis of Ran BP-1 expression in NIH/3T3 (A) and F9 (B) whole cell lysates.

SELECT PRODUCT CITATIONS

- Liu, X., et al. 2014. CRM1 is a direct cellular target of the natural anti-cancer agent plumbagin. *J. Pharmacol. Sci.* 124: 486-493.
- Munnur, D., et al. 2019. NR4A nuclear receptors target poly-ADP ribosylated DNA-PK_{CS} protein to promote DNA repair. *Cell Rep.* 26: 2028-2036.e6.
- Gao, J., et al. 2021. Discovery and biological evaluation of a small-molecule inhibitor of CRM1 that suppresses the growth of triple-negative breast cancer cells. *Traffic* 22: 221-229.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA