

Fc ϵ R γ (H-5): sc-374385

BACKGROUND

IgE Fc receptor I binds to the Fc region of immunoglobulins ϵ chain with high affinity, and is responsible for initiating the allergic response. Binding of allergen to receptor-bound IgE leads to cell activation and the release of mediators such as histamines, responsible for the manifestations of allergy. IgE Fc receptor I also induces the secretion of important lymphokines, effectors of the hypersensitivity response. It is a tetramer of a heavily glycosylated α chain, a β chain, and two disulfide linked γ chains. The γ chains from Fc ϵ RI are also subunits of other Fc receptors. The γ subunit is thought to be functionally significant in allowing the IgE Fc receptor to reach the cell surface. The cytoplasmic domains of the β and γ subunits each contain a conserved consensus sequence, ITAM, (immunoreceptor tyrosine activation motif). Phosphorylation of a pair of conserved tyrosine residues within this motif is required for signal transduction in mast cells and other hemopoietic cell types.

REFERENCES

- Hackel, W., et al. 1968. Foreign body as cause of a large urethral calculus and diverticulum formation. *Z. Urol. Nephrol.* 61: 827-829.
- Shimizu, A., et al. 1988. Human and rat mast cell high-affinity immunoglobulin E receptors: characterization of putative α -chain gene products. *Proc. Natl. Acad. Sci. USA* 85: 1907-1911.
- Le Coniat, M., et al. 1990. The human genes for the α and γ subunits of the mast cell receptor for immunoglobulin E are located on human chromosome band 1q23. *Immunogenetics* 32: 183-186.
- Kuster, H., et al. 1992. The gene and cDNA for the human high affinity immunoglobulin E receptor β chain and expression of the complete human receptor. *J. Biol. Chem.* 267: 12782-12787.
- Pang, J., et al. 1993. Characterization of the gene for the human high affinity IgE receptor (Fc ϵ RI) α -chain. *J. Immunol.* 151: 6166-6174.
- Penhallow, R.C., et al. 1995. Temporal activation of nontransmembrane protein-tyrosine kinases following mast cell Fc ϵ RI engagement. *J. Biol. Chem.* 270: 23362-23365.

CHROMOSOMAL LOCATION

Genetic locus: FCER1G (human) mapping to 1q23.3; Fc ϵ r1g (mouse) mapping to 1 H3.

SOURCE

Fc ϵ R γ (H-5) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 51-85 within a C-terminal cytoplasmic domain of Fc ϵ R γ of human origin.

PRODUCT

Each vial contains 200 μ g IgG $_1$ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-374385 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

Fc ϵ R γ (H-5) is recommended for detection of Fc ϵ R γ of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Fc ϵ R γ (H-5) is also recommended for detection of Fc ϵ R γ in additional species, including porcine.

Suitable for use as control antibody for Fc ϵ R γ siRNA (h): sc-45267, Fc ϵ R γ siRNA (m): sc-45268, Fc ϵ R γ shRNA Plasmid (h): sc-45267-SH, Fc ϵ R γ shRNA Plasmid (m): sc-45268-SH, Fc ϵ R γ shRNA (h) Lentiviral Particles: sc-45267-V and Fc ϵ R γ shRNA (m) Lentiviral Particles: sc-45268-V.


Molecular Weight of Fc ϵ R γ : 9 kDa.

Positive Controls: CTLL-2 cell lysate: sc-2242, THP-1 cell lysate: sc-2238, Fc ϵ R γ (h): 293T Lysate: sc-115131.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



Fc ϵ R γ (H-5): sc-374385. Western blot analysis of Fc ϵ R γ expression in non-transfected: sc-117752 (A) and human Fc ϵ R γ transfected: sc-115131 (B) 293T whole cell lysates.



Fc ϵ R γ (H-5): sc-374385. Western blot analysis of Fc ϵ R γ expression in THP-1 whole cell lysate.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.