

# Aciculin (F-11): sc-374530

## BACKGROUND

Aciculin, also known as PGM5 (phosphoglucomutase 5) or PGMRP, is a 567 amino acid protein that belongs to the phosphoglucomutase family of phosphotransferases, which play an important role in the interconversion of glucose-1-phosphate and glucose-6-phosphate. Localized to the cell junction and expressed at high levels in smooth and cardiac muscle, Aciculin binds magnesium as a cofactor and interacts with dystrophin and utrophin, possibly playing a role in cytoskeletal organization and function. Aciculin exists as two alternatively spliced isoforms and is encoded by a gene that maps to human chromosome 9, which houses over 900 genes and comprises nearly 4% of the human genome.

## REFERENCES

1. Belkin, A.M., et al. 1994. Expression and localization of the phosphoglucomutase-related cytoskeletal protein, Aciculin, in skeletal muscle. *J. Cell Sci.* 107: 1993-2003.
2. Belkin, A.M., et al. 1995. Localization of utrophin and Aciculin at sites of cell-matrix and cell-cell adhesion in cultured cells. *Exp. Cell Res.* 221: 132-140.
3. Belkin, A.M., et al. 1995. Association of Aciculin with dystrophin and utrophin. *J. Biol. Chem.* 270: 6328-6337.
4. Belkin, A.M., et al. 1996. Localization of crinin (dystroglycan) at sites of cell-matrix and cell-cell contact: recruitment to focal adhesions is dependent upon extracellular ligands. *Cell Adhes. Commun.* 4: 281-296.
5. Moiseeva, E.P., et al. 1996. A novel dystrophin/utrophin-associated protein is an enzymatically inactive member of the phosphoglucomutase superfamily. *Eur. J. Biochem.* 235: 103-113.
6. Rezvani, M., et al. 1996. Dystrophin, vinculin, and Aciculin in skeletal muscle subject to chronic use and disuse. *Med. Sci. Sports Exerc.* 28: 79-84.
7. Moiseeva, E.P., et al. 1997. Characterisation of the promoter which regulates expression of a phosphoglucomutase-related protein, a component of the dystrophin/utrophin cytoskeleton predominantly expressed in smooth muscle. *Eur. J. Biochem.* 248: 634-643.
8. Wakayama, Y., et al. 2000. Aciculin and its relation to dystrophin: immunocytochemical studies in human normal and Duchenne dystrophy quadriceps muscles. *Acta Neuropathol.* 99: 654-662.
9. Online Mendelian Inheritance in Man, OMIM™. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 600981. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

## CHROMOSOMAL LOCATION

Genetic locus: PGM5 (human) mapping to 9q21.11; Pgm5 (mouse) mapping to 19 B.

## SOURCE

Aciculin (F-11) is a mouse monoclonal antibody raised against amino acids 428-484 mapping near the C-terminus of Aciculin of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>2a</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

Aciculin (F-11) is recommended for detection of Aciculin of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Aciculin siRNA (h): sc-72431, Aciculin siRNA (m): sc-72432, Aciculin shRNA Plasmid (h): sc-72431-SH, Aciculin shRNA Plasmid (m): sc-72432-SH, Aciculin shRNA (h) Lentiviral Particles: sc-72431-V and Aciculin shRNA (m) Lentiviral Particles: sc-72432-V.


Molecular Weight of Aciculin: 56 kDa.

Positive Controls: A-10 cell lysate: sc-3806, SK-BR-3 cell lysate: sc-2218 or human kidney extract: sc-363764.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



Aciculin (F-11): sc-374530. Western blot analysis of Aciculin expression in A-10 whole cell lysate (A) and human kidney tissue extract (B).



Aciculin (F-11): sc-374530. Western blot analysis of Aciculin expression in SK-BR-3 whole cell lysate.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.