

TAF II p100 (H-3): sc-374644

BACKGROUND

TFIID is a general transcription factor which initiates preinitiation complex assembly through direct interaction with the TATA promoter element. It is a multisubunit complex consisting of a small TATA-binding polypeptide and other TATA-binding protein (TBP)-associated factors (TAFs). Although native TFIID can mediate both activator-independent (basal) and activator-dependent transcription in reconstituted systems, TBP can mediate only basal transcription. TAF II p100 (TBP-associated factor II100), also known as TAF5 or TAFII100, is the third largest subunit of human TFIID. It contains six WD40 repeats at the C-terminus and has an N-terminus capable of forming a flexible dimer. TAF II p100 plays an important role in forming the scaffold that is crucial for the assembly of the TFIID complex. TAF II p100 may also be involved in the stabilization of TAF interactions.

REFERENCES

1. Matsui, T., et al. 1980. Multiple factors required for accurate initiation of transcription by purified RNA polymerase II. *J. Biol. Chem.* 255: 11992-11996.
2. Buratowski, S., et al. 1989. Five intermediate complexes in transcription initiation by RNA polymerase II. *Cell* 56: 549-561.
3. Takada, R., et al. 1992. Identification of human TFIID components and direct interaction between a 250 kDa polypeptide and the TATA box-binding protein (TFIIDt). *Proc. Natl. Acad. Sci. USA* 89: 11809-11813.
4. Bellorini, M., et al. 1997. CCAAT binding NF-Y-TBP interactions: NF-YB and NF-YC require short domains adjacent to their histone fold motifs for association with TBP basic residues. *Nucleic Acids Res.* 25: 2174-2181.
5. Tao, Y., et al. 1997. Specific interactions and potential functions of human TAFII100. *J. Biol. Chem.* 272: 6714-6721.

CHROMOSOMAL LOCATION

Genetic locus: TAF5 (human) mapping to 10q24.33; Taf5 (mouse) mapping to 19 C3.

SOURCE

TAF II p100 (H-3) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 263-293 within an internal region of TAF II p100 of human origin.

PRODUCT

Each vial contains 200 µg IgG₃ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-374644 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

TAF II p100 (H-3) is recommended for detection of TAF II p100 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

TAF II p100 (H-3) is also recommended for detection of TAF II p100 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for TAF II p100 siRNA (h): sc-38498, TAF II p100 siRNA (m): sc-154044, TAF II p100 shRNA Plasmid (h): sc-38498-SH, TAF II p100 shRNA Plasmid (m): sc-154044-SH, TAF II p100 shRNA (h) Lentiviral Particles: sc-38498-V and TAF II p100 shRNA (m) Lentiviral Particles: sc-154044-V.

Molecular Weight (predicted) of short/long TAF II p100 isoforms: 81/87 kDa.


Molecular Weight (observed) of TAF II p100: 80-101 kDa.

Positive Controls: MCF7 whole cell lysate: sc-2206, HeLa nuclear extract: sc-2120 or HeLa whole cell lysate: sc-2200.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



TAF II p100 (H-3): sc-374644. Western blot analysis of TAF II p100 expression in HeLa (A) and MCF7 (B) whole cell lysates.



TAF II p100 (H-3): sc-374644. Immunoperoxidase staining of formalin fixed, paraffin-embedded human adrenal gland tissue showing nuclear and cytoplasmic staining of glandular cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.