FAM123A (G-1): sc-374654



The Power to Ouestion

BACKGROUND

The FAM123A (family with sequence similarity 123A) gene encodes for a 671 amino acid protein. There are two isoforms of FAM123A that exist as a result of alternative splicing events. The gene encoding FAM123A is located on chromosome 13, which comprises nearly 4% of human DNA and contains about 114 million base pairs and 400 genes. Key tumor suppressor genes on chromosome 13 include the breast cancer susceptibility gene, BRCA2, and the RB1 (retinoblastoma) gene. RB1 encodes a crucial tumor suppressor protein which, when defective, leads to malignant growth in the retina and has been implicated in a variety of other cancers. The gene SLITRK1, which is associated with Tourette syndrome, is on chromosome 13. As with most chromosomes, polysomy of part or all of chromosome 13 is deleterious to development and decreases the odds of survival.

REFERENCES

- Dunham, A., et al. 2004. The DNA sequence and analysis of human chromosome 13. Nature 428: 522-528.
- Deng, H., et al. 2006. Examination of the SLITRK1 gene in Caucasian patients with Tourette syndrome. Acta Neurol. Scand. 114: 400-402.
- 3. Giacinti, C., et al. 2006. RB and cell cycle progression. Oncogene 25: 5220-5227.
- 4. Grados, M.A., et al. 2006. A new gene for Tourette's syndrome: a window into causal mechanisms? Trends Genet. 22: 291-293.
- Bugge, M., et al. 2007. Non-disjunction of chromosome 13. Hum. Mol. Genet. 16: 2004-2010.

CHROMOSOMAL LOCATION

Genetic locus: AMER2 (human) mapping to 13q12.13; Amer2 (mouse) mapping to 14 D1.

SOURCE

FAM123A (G-1) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 603-637 near the C-terminus of FAM123A of human origin.

PRODUCT

Each vial contains 200 μg lgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

FAM123A (G-1) is available conjugated to agarose (sc-374654 AC), 500 μg/ 0.25 ml agarose in 1 ml, for IP; to HRP (sc-374654 HRP), 200 μg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-374654 PE), fluorescein (sc-374654 FITC), Alexa Fluor® 488 (sc-374654 AF488), Alexa Fluor® 546 (sc-374654 AF546), Alexa Fluor® 594 (sc-374654 AF594) or Alexa Fluor® 647 (sc-374654 AF647), 200 μg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-374654 AF680) or Alexa Fluor® 790 (sc-374654 AF790), 200 μg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-374654 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

FAM123A (G-1) is recommended for detection of FAM123A of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for FAM123A siRNA (h): sc-105346, FAM123A siRNA (m): sc-108774, FAM123A shRNA Plasmid (h): sc-105346-SH, FAM123A shRNA Plasmid (m): sc-108774-SH, FAM123A shRNA (h) Lentiviral Particles: sc-105346-V and FAM123A shRNA (m) Lentiviral Particles: sc-108774-V.


Molecular Weight of FAM123A: 70/58 kDa.


Positive Controls: FAM123A (h): 293T Lysate: sc-115267.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-lgG κ BP-HRP: sc-516102 or m-lgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz MarkerTM Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-lgG κ BP-FITC: sc-516140 or m-lgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA







FAM123A (G-1): sc-374654. Immunofluorescence staining of methanol-fixed HeLa cells showing nuclear and cytoplasmic localization.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA