

FAM123A (G-1): sc-374654

BACKGROUND

The FAM123A (family with sequence similarity 123A) gene encodes for a 671 amino acid protein. There are two isoforms of FAM123A that exist as a result of alternative splicing events. The gene encoding FAM123A is located on chromosome 13, which comprises nearly 4% of human DNA and contains about 114 million base pairs and 400 genes. Key tumor suppressor genes on chromosome 13 include the breast cancer susceptibility gene, BRCA2, and the RB1 (retinoblastoma) gene. RB1 encodes a crucial tumor suppressor protein which, when defective, leads to malignant growth in the retina and has been implicated in a variety of other cancers. The gene SLITRK1, which is associated with Tourette syndrome, is on chromosome 13. As with most chromosomes, polysomy of part or all of chromosome 13 is deleterious to development and decreases the odds of survival.

REFERENCES

1. Dunham, A., et al. 2004. The DNA sequence and analysis of human chromosome 13. *Nature* 428: 522-528.
2. Deng, H., et al. 2006. Examination of the SLITRK1 gene in Caucasian patients with Tourette syndrome. *Acta Neurol. Scand.* 114: 400-402.
3. Giacinti, C., et al. 2006. RB and cell cycle progression. *Oncogene* 25: 5220-5227.
4. Grados, M.A., et al. 2006. A new gene for Tourette's syndrome: a window into causal mechanisms? *Trends Genet.* 22: 291-293.
5. Bugge, M., et al. 2007. Non-disjunction of chromosome 13. *Hum. Mol. Genet.* 16: 2004-2010.

CHROMOSOMAL LOCATION

Genetic locus: AMER2 (human) mapping to 13q12.13; Amer2 (mouse) mapping to 14 D1.

SOURCE

FAM123A (G-1) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 603-637 near the C-terminus of FAM123A of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

FAM123A (G-1) is available conjugated to agarose (sc-374654 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-374654 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-374654 PE), fluorescein (sc-374654 FITC), Alexa Fluor® 488 (sc-374654 AF488), Alexa Fluor® 546 (sc-374654 AF546), Alexa Fluor® 594 (sc-374654 AF594) or Alexa Fluor® 647 (sc-374654 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-374654 AF680) or Alexa Fluor® 790 (sc-374654 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-374654 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

FAM123A (G-1) is recommended for detection of FAM123A of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for FAM123A siRNA (h): sc-105346, FAM123A siRNA (m): sc-108774, FAM123A shRNA Plasmid (h): sc-105346-SH, FAM123A shRNA Plasmid (m): sc-108774-SH, FAM123A shRNA (h) Lentiviral Particles: sc-105346-V and FAM123A shRNA (m) Lentiviral Particles: sc-108774-V.

Molecular Weight of FAM123A: 70/58 kDa.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



FAM123A (G-1): sc-374654. Western blot analysis of FAM123A expression in untreated (A) and chemically-treated (B, C, D) whole cell lysates. Detection reagent used: m-IgGκ BP-HRP: sc-516102. β-Actin (C4): sc-47778 used as loading control. Detection reagent used: m-IgG Fc BP-HRP: sc-525409.



FAM123A (G-1): sc-374654. Immunofluorescence staining of methanol-fixed HeLa cells showing nuclear and cytoplasmic localization.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA