SANTA CRUZ BIOTECHNOLOGY, INC.

PKC v (C-1): sc-376024



BACKGROUND

Members of the protein kinase C (PKC) family play a key regulatory role in a variety of cellular functions including cell growth and differentiation, gene expression, hormone secretion and membrane function. PKCs were originally identified as serine/threonine protein kinases whose activity was dependent on calcium and phospholipids. Diacylglycerols (DAG) and tumor promoting phorbol esters bind to and activate PKC. PKCs can be subdivided into at least two major classes, including conventional (c) PKC isoforms (α , β I, β II, γ) and novel (n) PKC isoforms (δ , ε , ζ , η , θ , λ/ι , μ and ν). Patterns of expression for each PKC offerences in their cofactor dependencies. For instance, the kinase activities of PKC δ and ε are independent of Ca²⁺. On the other hand, most of the other PKC members possess phorbol ester-binding activities and kinase activities.

REFERENCES

- Takai, Y., et al. 1979. Calcium-dependent activation of a multifunctional protein kinase by membrane phospholipids. J. Biol. Chem. 254: 3692-3695.
- Castagna, M., et al. 1982. Direct activation of calcium-activated, phospholipid-dependent protein kinase by tumor-promoting phorbol esters. J. Biol. Chem. 257: 7847-7851.
- Kikkawa, U., et al. 1983. Protein kinase C as a possible receptor of tumorpromoting phorbol esters. J. Biol. Chem. 258: 11442-11445.

CHROMOSOMAL LOCATION

Genetic locus: PRKD3 (human) mapping to 2p22.2; Prkd3 (mouse) mapping to 17 E3.

SOURCE

PKC v (C-1) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 31-69 near the N-terminus of PKC v of human origin.

PRODUCT

Each vial contains 200 $\mu g\, lgG_{2b}$ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

PKC v (C-1) is available conjugated to agarose (sc-376024 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-376024 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-376024 PE), fluorescein (sc-376024 FITC), Alexa Fluor[®] 488 (sc-376024 AF488), Alexa Fluor[®] 546 (sc-376024 AF546), Alexa Fluor[®] 594 (sc-376024 AF594) or Alexa Fluor[®] 647 (sc-376024 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-376024 AF680) or Alexa Fluor[®] 790 (sc-376024 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-376024 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

PKC v (C-1) is recommended for detection of PKC v of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

PKC ν (C-1) is also recommended for detection of PKC ν in additional species, including bovine.

Suitable for use as control antibody for PKC ν siRNA (h): sc-44789, PKC ν siRNA (m): sc-44790, PKC ν shRNA Plasmid (h): sc-44789-SH, PKC ν shRNA Plasmid (m): sc-44790-SH, PKC ν shRNA (h) Lentiviral Particles: sc-44789-V and PKC ν shRNA (m) Lentiviral Particles: sc-44790-V.

Molecular Weight of PKC v: 100 kDa.

Positive Controls: PKC ν (h): 293 Lysate: sc-158862, Ramos cell lysate: sc-2216 or HeLa whole cell lysate: sc-2200.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA





 $\mathsf{PKC}\,\nu$ (C-1): sc-376024. Western blot analysis of $\mathsf{PKC}\,\nu$ expression in non-transfected: sc-110760 (A) and human $\mathsf{PKC}\,\nu$ transfected: sc-158862 (B) 293 whole cell lysates.

PKC v (C-1): sc-376024. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic and membrane localization (A). Immunoperoxidaes staining of formalin fixed, paraffin-embedded human kidney tissue showing cytoplasmic and apical membrane staining of cells in tubules (B).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.