# PQBP-1 (G-12): sc-376039



The Power to Question

### **BACKGROUND**

Polyglutamine(Q) tract binding protein-1 (PQBP-1) is a transcription repressor that associates with polyglutamine tract-containing transcription regulators and causative genes for neurodegenerative disorders. Hepta- and di-amino acid repeat sequences rich in polar residues are essential for PQBP-1 to interact with polyglutamine tract-containing proteins (i.e. huntingtin, androgen receptor and Brain-2). PQBP-1 contains a WWP/WW domain that binds proline-rich motifs and a C2 domain that can influence Ca<sup>2+</sup>-dependent phospholipid signaling. PQBP-1 localizes to the nucleus and is present in neurons throughout the brain, with abundant levels in hippocampus, cerebellar cortex and olfactory bulb. The human PQBP-1 gene maps to chromosome Xp11.23.

#### **REFERENCES**

- Imafuku, I., et al. 1998. Polar amino acid-rich sequences bind to polyglutamine tracts. Biochem. Biophys. Res. Commun. 253: 16-20.
- 2. Waragai, M., et al. 1999. POBP-1, a novel polyglutamine tract-binding protein, inhibits transcription activation by Brn-2 and affects cell survival. Hum. Mol. Genet. 8: 977-987.

### **CHROMOSOMAL LOCATION**

Genetic locus: PQBP1 (human) mapping to Xp11.23; Pqbp1 (mouse) mapping to X A1.1.

#### **SOURCE**

PQBP-1 (G-12) is a mouse monoclonal antibody raised against amino acids 1-265 representing full length PQBP-1 of human origin.

### **PRODUCT**

Each vial contains 200  $\mu$ g lgG<sub>2b</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-376039 X, 200  $\mu$ g/0.1 ml.

## **APPLICATIONS**

POBP-1 (G-12) is recommended for detection of POBP-1 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for PQBP-1 siRNA (h): sc-38199, PQBP-1 siRNA (m): sc-38200, PQBP-1 shRNA Plasmid (h): sc-38199-SH, PQBP-1 shRNA Plasmid (m): sc-38200-SH, PQBP-1 shRNA (h) Lentiviral Particles: sc-38199-V and PQBP-1 shRNA (m) Lentiviral Particles: sc-38200-V.

PQBP-1 (G-12) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of PQBP-1: 38 kDa.


Positive Controls: Sol8 nuclear extract: sc-2157 or PQBP-1 (m): 293T Lysate: sc-122739.

### **RECOMMENDED SUPPORT REAGENTS**

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-lgG $\kappa$  BP-HRP: sc-516102 or m-lgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>TM</sup> Molecular Weight Standards: sc-2035, UltraCruz\* Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-lgG $\kappa$  BP-FITC: sc-516140 or m-lgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz\* Mounting Medium: sc-24941 or UltraCruz\* Hard-set Mounting Medium: sc-359850.

#### **DATA**





PQBP-1 (G-12): sc-376039. Western blot analysis of PQBP-1 expression in non-transfected: sc-117752 (A) and mouse PQBP-1 transfected: sc-122739 (B) 293T whole cell Ivsates.

PQBP-1 (G-12): sc-376039. Immunofluorescence staining of methanol-fixed HeLa cells showing nucleolar and cytoplasmic localization.

#### **SELECT PRODUCT CITATIONS**

- Fuchs, N.V., et al. 2019. Induced pluripotent stem cells (iPSCs) derived from a renpenning syndrome patient with c.459\_462delAGAG mutation in POBP1 (PEli001-A). Stem Cell Res. 41: 101592.
- 2. Yoh, S.M., et al. 2022. Recognition of HIV-1 capsid by PQBP1 licenses an innate immune sensing of nascent HIV-1 DNA. Mol. Cell 82: 2871-2884.e6.

# **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

# **PROTOCOLS**

See our web site at www.scbt.com for detailed protocols and support products.