# SANTA CRUZ BIOTECHNOLOGY, INC.

# UFSP2 (C-7): sc-376045



#### BACKGROUND

UFM1 (ubiquitin-fold modifier 1) is a ubiquitin-like protein that is conjugated to target proteins by UBA5, an E1-like activating enzyme, and Ufc1, an E2-like conjugating enzyme. Through these interactions, UFM1 conjugates to target proteins by a covalent linkage. UFSP2 (Ufm1-specific protease 2) is a 469 amino acid thiol protease that specifically cleaves UFM1 precursor and leads to exposure of its conserved C-terminal glycine, a step required prior to conjugation to target proteins. UFSP2 is also capable of releasing UFM1 from UFM1-conjugated cellular proteins. The gene encoding UFSP2 maps to human chromosome 4, which houses nearly 6% of the human genome and has the largest gene deserts (regions of the genome with no protein encoding genes) of all of the human chromosomes. Defects in some of the genes located on chromosome 4 are associated with Huntington's disease, Ellis-van Creveld syndrome, methylmalonic acidemia and polycystic kidney disease.

#### REFERENCES

- 1. Tanaka, K., et al. 1998. The ligation systems for ubiquitin and ubiquitin-like proteins. Mol. Cells 8: 503-512.
- Wilson, M.D., et al. 2001. Comparative analysis of the gene-dense ACHE/ TFR2 region on human chromosome 7q22 with the orthologous region on mouse chromosome 5. Nucleic Acids Res. 29: 1352-1365.
- Komatsu, M., et al. 2004. A novel protein-conjugating system for Ufm1, a ubiquitin-fold modifier. EMBO J. 23: 1977-1986.
- Kang, S.H., et al. 2007. Two novel ubiquitin-fold modifier 1 (Ufm1)-specific proteases, UFSP1 and UFSP2. J. Biol. Chem. 282: 5256-5262.
- 5. Ha, B.H., et al. 2008. Structural basis for Ufm1 processing by UFSP1. J. Biol. Chem. 283: 14893-14900.
- 6. Online Mendelian Inheritance in Man, OMIM<sup>™</sup>. 2009. Johns Hopkins University, Baltimore, MD. MIM Number: 611482. World Wide Web URL: http://www.ncbi.nlm.nih.gov/omim/
- Eijgelsheim, M., et al. 2010. Genome-wide association analysis identifies multiple loci related to resting heart rate. Hum. Mol. Genet. 19: 3885-3894.
- Tatsumi, K., et al. 2010. A novel type of E3 ligase for the Ufm1 conjugation system. J. Biol. Chem. 285: 5417-5427.

#### **CHROMOSOMAL LOCATION**

Genetic locus: UFSP2 (human) mapping to 4q35.1; Ufsp2 (mouse) mapping to 8 B1.1.

## SOURCE

UFSP2 (C-7) is a mouse monoclonal antibody raised against amino acids 155-340 mapping within an internal region of UFSP2 of human origin.

## PRODUCT

Each vial contains 200  $\mu g$  lgG\_{2b} lambda light chain in 1.0 ml of PBS with <0.1% sodium azide and 0.1% gelatin.

#### **APPLICATIONS**

UFSP2 (C-7) is recommended for detection of UFSP2 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for UFSP2 siRNA (h): sc-89115, UFSP2 siRNA (m): sc-154893, UFSP2 shRNA Plasmid (h): sc-89115-SH, UFSP2 shRNA Plasmid (m): sc-154893-SH, UFSP2 shRNA (h) Lentiviral Particles: sc-89115-V and UFSP2 shRNA (m) Lentiviral Particles: sc-154893-V.

Molecular Weight of UFSP2: 53 kDa.

Positive Controls: Caki-1 cell lysate: sc-2224, TF-1 cell lysate: sc-2412 or U-87 MG cell lysate: sc-2411.

## **RECOMMENDED SUPPORT REAGENTS**

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\lambda$  BP-HRP: sc-516132 or m-IgG $\lambda$  BP-HRP (Cruz Marker): sc-516132-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>TM</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG $\lambda$  BP-FITC: sc-516185 or m-IgG $\lambda$  BP-PE: sc-516186 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG $\lambda$  BP-HRP: sc-516132 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

## DATA





UFSP2 (C-7): sc-376045. Western blot analysis of UFSP2 expression in Caki-1 (A), U-87 MG (B) and TF-1 (C) whole cell lysates. Detection reagent used: m-IgG $\lambda$  BP-HRP (Cruz Marker): sc-516132-CM.

UFSP2 (C-7): sc-376045. Immunofluorescence staining of methanol-fixed HeLa cells showing nuclear and cytoplasmic localization (**A**). Immunoperoxidase staining of formalin fixed, paraffin-embedded human adrenal gland tissue showing cytoplasmic staining of glandular cells (**B**).

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## **RESEARCH USE**

For research use only, not for use in diagnostic procedures.