SANTA CRUZ BIOTECHNOLOGY, INC.

γ-parvin (B-9): sc-376081



BACKGROUND

The parvin family, including α -parvin, β -parvin and γ -parvin, link integrins and associated proteins with intracellular pathways, which regulate Actin cytoskeletal dynamics and cell survival. All three family members localize to focal adhesions and function in cell adhesion, spreading, motility and survival through interactions with partners, such as integrin-linked kinase (ILK), paxillin, α -actinin and testicular kinase 1. α -parvin is widely expressed, with highest levels detected in the skeletal muscle, heart, liver and kidney. A complex made up of α -parvin, ILK and the LIM protein PINCH-1 is critical for cell survival in a variety of cells, including certain cancer cells, kidney podocytes and cardiac myocytes. β -parvin links initial integrin signals to rapid Actin reorganization, thereby playing a critical role in fibroblast migration. The ILK- γ -parvin complex is essential for the establishment of cell polarity required for leukocyte migration.

REFERENCES

- 1. Olski, T.M., et al. 2001. Parvin, a 42 kDa focal adhesion protein, related to the α -actinin superfamily. J. Cell Sci. 114: 525-538.
- 2. Korenbaum, E., et al. 2001. Genomic organization and expression profile of the parvin family of focal adhesion proteins in mice and humans. Gene 279: 69-79.
- Aboulaich, N., et al. 2004. Vectorial proteomics reveal targeting, of polymerase I and transcript release factor (PTRF) at the surface of caveolae in human adipocytes. Biochem. J. 383: 237-248.
- 4. Yamaji, S., et al. 2004. Affixin interacts with α -actinin and mediates integrin signaling for reorganization of F-Actin induced by initial cell-substrate interaction. J. Cell Biol. 165: 539-551.
- 5. Zhang, Y., et al. 2004. Distinct roles of two structurally closely related focal adhesion proteins, α -parvins and β -parvins, in regulation of cell morphology and survival. J. Biol. Chem. 279: 41695-41705.
- 6. Filipenko, N.R., et al. 2005. Integrin-linked kinase activity regulates Racand Cdc42-mediated Actin cytoskeleton reorganization via α -PIX. Oncogene 24: 5837-5849.

CHROMOSOMAL LOCATION

Genetic locus: PARVG (human) mapping to 22q13.31; Parvg (mouse) mapping to 15 E2.

SOURCE

 γ -parvin (B-9) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 41-79 near the N-terminus of γ -parvin of human origin.

PRODUCT

Each vial contains 200 μg IgG_1 kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-376081 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

 γ -parvin (B-9) is recommended for detection of γ -parvin of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for γ -parvin siRNA (h): sc-61302, γ -parvin siRNA (m): sc-61304, γ -parvin shRNA Plasmid (h): sc-61302-SH, γ -parvin shRNA Plasmid (m): sc-61304-SH, γ -parvin shRNA (h) Lentiviral Particles: sc-61302-V and γ -parvin shRNA (m) Lentiviral Particles: sc-61304-V.

Molecular Weight of y-parvin: 37 kDa.

Positive Controls: γ -parvin (h): 293T Lysate: sc-114463 or HL-60 whole cell lysate: sc-2209.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA





γ-parvin (B-9): sc-376081. Western blot analysis of γ-parvin expression in non-transfected: sc-117752 (**A**) and human γ-parvin transfected: sc-114463 (**B**) 293T whole cell lysates.

γ-parvin (B-9): sc-376081. Immunoperoxidase staining of formalin fixed, paraffin-embedded human thyroid gland tissue showing cytoplasmic staining of glandular cells.

STORAGE

Store at 4° C, **D0 NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.