

VASP (D-11): sc-376226



The Power to Question

BACKGROUND

The Wiskott-Aldrich syndrome (WAS) is characterized by thrombocytopenia, eczema, defects in cell-mediated and humoral immunity, and a propensity for lymphoproliferative diseases. The syndrome is the result of a mutation in the gene encoding a proline-rich protein termed WASP. WASP has been identified as a downstream effector of Cdc42 and has been implicated in Actin polymerization and cytoskeletal organization. A distantly related protein, VASP (vaso-dilator-stimulated phosphoprotein), is involved in the maintenance of cytoarchitecture by interacting with Actin-like filaments. VASP shares a limited degree of homology with the amino-terminus of WASP, which is frequently mutated in WAS patients. An established substrate of cAMP and cGMP dependent kinases, VASP is phosphorylated on a regulatory Serine residue 157 and localizes to focal adhesions, microfilaments and highly active regions of the plasma membrane. VASP is highly expressed in human platelets and, like WASP, may play a role in cytoskeletal organization.

CHROMOSOMAL LOCATION

Genetic locus: VASP (human) mapping to 19q13.32; Vasp (mouse) mapping to 7 A3.

SOURCE

VASP (D-11) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 353-385 near the C-terminus of VASP of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-376226 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

VASP (D-11) is recommended for detection of VASP of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

VASP (D-11) is also recommended for detection of VASP in additional species, including canine.

Suitable for use as control antibody for VASP siRNA (h): sc-29516, VASP siRNA (m): sc-36809, VASP shRNA Plasmid (h): sc-29516-SH, VASP shRNA Plasmid (m): sc-36809-SH, VASP shRNA (h) Lentiviral Particles: sc-29516-V and VASP shRNA (m) Lentiviral Particles: sc-36809-V.

Molecular Weight of VASP: 46 kDa.


Molecular Weight of phosphorylated VASP: 50 kDa.

Positive Controls: MTE1D whole cell lysate: sc-364918, A-431 whole cell lysate: sc-2201 or VASP (h): 293T Lysate: sc-114829.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



VASP (D-11): sc-376226. Western blot analysis of VASP expression in non-transfected: sc-117752 (A) and human VASP transfected: sc-114829 (B) 293T whole cell lysates.



VASP (D-11): sc-376226. Western blot analysis of VASP expression in MTE1D whole cell lysate.

SELECT PRODUCT CITATIONS

1. Sanchez-Ruiloba, L., et al. 2014. Protein kinase D interacts with neuronal nitric oxide synthase and phosphorylates the activatory residue serine 1412. PLoS ONE 9: 1-17.
2. Ma, Q., et al. 2019. Antitumor effects of saikosaponin b2 on breast cancer cell proliferation and migration. Mol. Med. Rep. 20: 1943-1951.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.