SANTA CRUZ BIOTECHNOLOGY, INC.

ARA54 (B-1): sc-376285



BACKGROUND

Androgens exhibit a wide range of effects on the development, maintenance and regulation of male phenotype and reproductive physiology in males. The androgen receptor (AR) is a member of the steroid superfamily of liganddependent transcription factors. ARs bind the two biologically active androgens, testosterone (T) and dihydrotestosterone (DHT). ARA54 contains a RING finger and functions as an AR coactivator. Testis, thymus, spleen, colon, prostate and uterus express ARA54 at high levels. ARA160 associates with an ATP-dependent chromatin remodeling factor known as the SNF/SWI complex. The FXXLF motif present in AR coactivators mediates their interaction with AR but not their transcription-related activity.

REFERENCES

- 1. Walsh, P.C., et al. 1974. Familial incomplete male pseudohermaphroditism type 2: decreased dihydro-testosterone formation in pseudovaginal perineoscrotal hypospadias. N. Engl. J. Med. 291: 944-949.
- Imperato-McGinley, J., et al. 1974. Steroid 5α-reductase deficiency in man: an inherited form of male pseudohermaphroditism. Science 186: 1213-1215.
- Garcia, J.A., et al. 1992. Cloning and chromosomal mapping of a human immunodeficiency virus 1 "TATA" element modulatory factor. Proc. Natl. Acad. Sci. USA 89: 9372-9376.
- 4. Zhou, Z.X., et al. 1994. The androgen receptor: an overview. Recent Prog. Horm. Res. 49: 249-274.
- Kang, H.Y., et al. 1999. Cloning and characterization of human prostate coactivator ARA54, a novel protein that associates with the androgen receptor. J. Biol. Chem. 274: 8570-8576.
- Ueki, N., et al. 1999. Isolation and characterization of a novel human gene (HFB30) which encodes a protein with a RING finger motif. Biochim. Biophys. Acta 1445: 232-236.
- 7. He, B., et al. 2002. The FXXLF motif mediates androgen receptor-specific interactions with coregulators. J. Biol. Chem. 277: 10226-10235.

CHROMOSOMAL LOCATION

Genetic locus: RNF14 (human) mapping to 5q31.3; Rnf14 (mouse) mapping to 18 B3.

SOURCE

ARA54 (B-1) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 423-459 near the C-terminus of ARA54 of human origin.

PRODUCT

Each vial contains 200 μ g lgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-376285 X, 200 μ g/0.1 ml.

Blocking peptide available for competition studies, sc-376285 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

ARA54 (B-1) is recommended for detection of ARA54 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for ARA54 siRNA (h): sc-43618, ARA54 siRNA (m): sc-44878, ARA54 shRNA Plasmid (h): sc-43618-SH, ARA54 shRNA Plasmid (m): sc-44878-SH, ARA54 shRNA (h) Lentiviral Particles: sc-43618-V and ARA54 shRNA (m) Lentiviral Particles: sc-44878-V.

ARA54 (B-1) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of ARA54: 54 kDa.

Positive Controls: DU 145 cell lysate: sc-2268.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

132 K -90 K -55 K -43 K -



ARA54 (B-1): sc-376285. Western blot analysis of ARA54 expression in DU 145 whole cell lysate.

ARA54 (B-1): sc-376285. Immunoperoxidase staining of formalin fixed, paraffin-embedded human testis tissue showing nuclear staining of subset of cells in seminiferous ducts.

STORAGE

Store at 4° C, **D0 NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.