

# PUS1 (C-4): sc-376442

## BACKGROUND

PUS1 (pseudouridine synthase 1) belongs to the tRNA pseudouridine synthase truA family. PUS1 functions in the conversion of uridine into pseudouridine after the nucleotide has been incorporated into RNA. It may have a functional role in tRNAs and is also thought to assist in the peptidyl transfer reaction of rRNAs. As a nucleus-resident protein, PUS1 forms a complex with RARG and the SRA1 RNA. PUS1 is widely expressed, with highest levels of expression in the brain and skeletal muscle tissues. Defects in PUS1 are a cause of myopathy with lactic acidosis and sideroblastic anemia (MLASA), also known as mitochondrial myopathy and sideroblastic anemia. MLASA is a rare autosomal recessive oxidative phosphorylation disorder specific to bone marrow and skeletal muscle. The deduced human PUS1 protein contains 348 amino acids and shares 92% sequence homology with mouse PUS1.

## REFERENCES

1. Arluison, V., et al. 1998. Transfer RNA-pseudouridine synthetase PUS1 of *Saccharomyces cerevisiae* contains one atom of zinc essential for its native conformation and tRNA recognition. *Biochemistry* 37: 7268-7276.
2. Arluison, V., et al. 1999. Pseudouridine synthetase Pus1 of *Saccharomyces cerevisiae*: kinetic characterisation, tRNA structural requirement and real-time analysis of its complex with tRNA. *J. Mol. Biol.* 289: 491-502.
3. Arluison, V., et al. 1999. RNA:pseudouridine synthetase Pus1 from *Saccharomyces cerevisiae*: oligomerization property and stoichiometry of the complex with yeast tRNA(Phe). *Biochimie* 81: 751-756.
4. Chen, J., et al. 1999. Cloning and characterization of a mammalian pseudouridine synthase. *RNA* 5: 409-419.
5. Chen, J., et al. 2000. Mouse pseudouridine synthase 1: gene structure and alternative splicing of pre-mRNA. *Biochem. J.* 352: 465-473.

## CHROMOSOMAL LOCATION

Genetic locus: PUS1 (human) mapping to 12q24.33; Pus1 (mouse) mapping to 5 F.

## SOURCE

PUS1 (C-4) is a mouse monoclonal antibody raised against amino acids 148-427 mapping at the C-terminus of PUS1 of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>2b</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

PUS1 (C-4) is available conjugated to agarose (sc-376442 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-376442 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-376442 PE), fluorescein (sc-376442 FITC), Alexa Fluor® 488 (sc-376442 AF488), Alexa Fluor® 546 (sc-376442 AF546), Alexa Fluor® 594 (sc-376442 AF594) or Alexa Fluor® 647 (sc-376442 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-376442 AF680) or Alexa Fluor® 790 (sc-376442 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

## APPLICATIONS

PUS1 (C-4) is recommended for detection of PUS1 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for PUS1 siRNA (h): sc-61417, PUS1 siRNA (m): sc-61418, PUS1 shRNA Plasmid (h): sc-61417-SH, PUS1 shRNA Plasmid (m): sc-61418-SH, PUS1 shRNA (h) Lentiviral Particles: sc-61417-V and PUS1 shRNA (m) Lentiviral Particles: sc-61418-V.


Molecular Weight of PUS1 isoform 1/2: 47/44 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, HeLa whole cell lysate: sc-2200 or A-431 whole cell lysate: sc-2201.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

## DATA



PUS1 (C-4): sc-376442. Western blot analysis of PUS1 expression in HeLa (A), A-431 (B) and Jurkat (C) whole cell lysates.



PUS1 (C-4): sc-376442. Immunofluorescence staining of formalin-fixed Hep G2 cells showing nuclear and cytoplasmic localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human upper stomach tissue showing nuclear and cytoplasmic staining of glandular cells (B).

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.