

nephrin (G-8): sc-376522

BACKGROUND

Nephrin is a member of the immunoglobulin family of cell adhesion molecules that localizes to opposing sites of the secondary foot processes formed by podocytes, a specialized epithelial cell that ensures size- and charge-selective ultrafiltration. The human nephrin gene maps to chromosome 19q13.12 and encodes a 1,241 amino acid protein that contains a transmembrane domain, eight Ig-like modules, and one fibronectin III-like module. Nephrin is expressed in embryonic and adult kidneys and localizes to glomerular podocytes and the glomerular slit diaphragm. Nephrin stimulates mitogen-activated protein kinases and is enhanced by podocin, which binds to the cytoplasmic tail of nephrin. A293 cells treated with phorbol-12-myristate-13-acetate can upregulate nephrin, suggesting that protein kinase C is part of an intracellular signalling system, which regulates nephrin.

CHROMOSOMAL LOCATION

Genetic locus: NPHS1 (human) mapping to 19q13.12; Nphs1 (mouse) mapping to 7 B1.

SOURCE

nephrin (G-8) is a mouse monoclonal antibody raised against amino acids 23-322 mapping within an extracellular domain of nephrin of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

nephrin (G-8) is available conjugated to agarose (sc-376522 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-376522 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-376522 PE), fluorescein (sc-376522 FITC), Alexa Fluor® 488 (sc-376522 AF488), Alexa Fluor® 546 (sc-376522 AF546), Alexa Fluor® 594 (sc-376522 AF594) or Alexa Fluor® 647 (sc-376522 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-376522 AF680) or Alexa Fluor® 790 (sc-376522 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

nephrin (G-8) is recommended for detection of nephrin of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for nephrin siRNA (h): sc-36030, nephrin siRNA (m): sc-36031, nephrin shRNA Plasmid (h): sc-36030-SH, nephrin shRNA Plasmid (m): sc-36031-SH, nephrin shRNA (h) Lentiviral Particles: sc-36030-V and nephrin shRNA (m) Lentiviral Particles: sc-36031-V.


Molecular Weight of nephrin: 185 kDa.

Positive Controls: human kidney extract: sc-363764, rat kidney extract: sc-2394 or KNRK whole cell lysate: sc-2214.


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA



nephrin (G-8): sc-376522. Western blot analysis of nephrin expression in human kidney tissue extract.



nephrin (G-8): sc-376522. Western blot analysis of nephrin expression in rat kidney tissue extract.

SELECT PRODUCT CITATIONS

- Cooper, C.J., et al. 2018. Characterization of a novel disease-associated mutation within NPHS1 and its effects on nephrin phosphorylation and signaling. *PLoS ONE* 13: e0203905.
- Nakuluri, K., et al. 2019. Hypoxia induces ZEB2 in podocytes: implications in the pathogenesis of proteinuria. *J. Cell. Physiol.* 234: 6503-6518.
- Chen, H., et al. 2020. Prenatal ethanol exposure increased the susceptibility of adult offspring rats to glomerulosclerosis. *Toxicol. Lett.* 321: 44-53.
- Bucur, O., et al. 2020. Nanoscale imaging of clinical specimens using conventional and rapid-expansion pathology. *Nat. Protoc.* 15: 1649-1672.
- Zhang, D., et al. 2020. *In vitro* induction and *in vivo* engraftment of kidney organoids derived from human pluripotent stem cells. *Exp. Ther. Med.* 20: 1307-1314.
- de Melo Junior, A.F., et al. 2020. Involvement of sex hormones, oxidative stress, ACE and ACE2 activity in the impairment of renal function and remodelling in SHR. *Life Sci.* 257: 118138.
- Luo, R., et al. 2020. A (pro)renin receptor decoy peptide PRO20 protects against adriamycin-induced nephropathy by targeting intrarenal renin-angiotensin system. *Am. J. Physiol. Renal Physiol.* 319: F930-F940.
- Maruyama-Fumoto, K., et al. 2021. Activation of protease-activated receptor 2 is associated with blood pressure regulation and proteinuria reduction in metabolic syndrome. *Clin. Exp. Pharmacol. Physiol.* 48: 211-220.
- Yamada, N., et al. 2021. Morphological analyses of nephrin expression in progressive glomerulonephropathy of common marmosets. *J. Toxicol. Pathol.* 34: 83-88.
- Chen, L., et al. 2021. Yiqi Huoxue recipe regulates autophagy through degradation of advanced glycation end products via mTOR/S6K1/LC3 pathway in diabetic nephropathy. *Evid. Based Complement. Alternat. Med.* 2021: 9942678.

RESEARCH USE

For research use only, not for use in diagnostic procedures.