

TRIP6 (G-5): sc-376550

BACKGROUND

Zyxin is a LIM domain-containing, zinc finger domain-containing, SH3 ligand-containing phosphoprotein that localizes to focal adhesion plaques and Actin filament bundles. Thyroid receptor interacting protein 6 (TRIP6) is a Zyxin-related protein. TRIP6 interacts with the ligand binding domain of the thyroid receptor and is predominantly expressed in kidney, liver and lung. It interacts with receptor-interacting protein 2 (RIP2) through LIM domains in a TNF- or IL-1-dependent manner. TRIP6 also interacts with TRAF2, a protein that is crucially involved in TNF signaling, as well as the IL-1 receptor, TLR2 and Nod1. Overexpression of TRIP6 facilitates NFκB activation by TNF, IL-1, TLR2 or Nod1, whereas a dominant negative mutant or RNA-interference construct of TRIP6 inhibits NFκB activation by TNF, IL-1, TLR2 or Nod1. Moreover, TRIP6 also potentiates RIP2- and Nod1-mediated ERK activation.

REFERENCES

1. Xu, J., et al. 2004. TRIP6 enhances lysophosphatidic acid-induced cell migration by interacting with the lysophosphatidic acid 2 receptor. *J. Biol. Chem.* 279: 10459-10468.
2. Lai, Y.J., et al. 2005. c-Src-mediated phosphorylation of TRIP6 regulates its function in lysophosphatidic acid-induced cell migration. *Mol. Cell. Biol.* 25: 5859-5868.
3. Li, L., et al. 2005. TRIP6 is a RIP2-associated common signaling component of multiple NFκB activation pathways. *J. Cell Sci.* 118: 555-563.
4. Petit, M.M., et al. 2005. The tumor suppressor Scrib selectively interacts with specific members of the Zyxin family of proteins. *FEBS Lett.* 579: 5061-5068.
5. Gur'ianova O.A., et al. 2005. Downregulation of TRIP6 expression induces Actin cytoskeleton rearrangements in human carcinoma cell lines. *Mol. Biol.* 39: 905-909.

CHROMOSOMAL LOCATION

Genetic locus: TRIP6 (human) mapping to 7q22.1; Trip6 (mouse) mapping to 5 G2.

SOURCE

TRIP6 (G-5) is a mouse monoclonal antibody raised against amino acids 160-254 mapping within an internal region of TRIP6 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

TRIP6 (G-5) is recommended for detection of TRIP6 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for TRIP6 siRNA (h): sc-45561, TRIP6 siRNA (m): sc-45562, TRIP6 shRNA Plasmid (h): sc-45561-SH, TRIP6 shRNA Plasmid (m): sc-45562-SH, TRIP6 shRNA (h) Lentiviral Particles: sc-45561-V and TRIP6 shRNA (m) Lentiviral Particles: sc-45562-V.

Molecular Weight of TRIP6: 50 kDa.


Positive Controls: HeLa nuclear extract: sc-2120, K-562 nuclear extract: sc-2130 or TRIP6 (h): 293T Lysate: sc-114028.

RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

- 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.
- 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).
- 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.
- 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



TRIP6 (G-5): sc-376550. Western blot analysis of TRIP6 expression in non-transfected: sc-117752 (A) and human TRIP6 transfected: sc-114028 (B) 293T whole cell lysates.



TRIP6 (G-5): sc-376550. Immunoperoxidase staining of formalin fixed, paraffin-embedded human uterine cervix tissue showing cytoplasmic and nuclear staining of squamous epithelial cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.