

ACAP1 (B-11): sc-376574

BACKGROUND

ACAP1, also designated Centaurin- β 1 (CENTB1 or Cnt-b1), is a member of the ADP ribosylation factor family of Arf6 GTPase-activating proteins (GAP). GAPs are important regulators of Arf function by controlling the return of Arf to its inactive state. ACAP1 is related to AGAP1 and ASAP1, and all three proteins are similarly expressed in fibroblast cells such as NIH/3T3. Internalization and recycling of integrin receptors is important in cell adhesion and migration modulation, and once inside a cell, proteins and membranes are transported to the endosome where they are sorted for recycling or degradation. ACAP1 promotes cargo sorting by associating directly to recycling cargo proteins. Preventing this interaction inhibits protein recycling. ACAP1 binds transferrin receptors, promoting their transport to the plasma membrane from the endosome. Akt induced phosphorylation of ACAP1 at Ser 554 regulates ACAP1 interaction to integrin in endosomes, and downregulation of Akt or ACAP1 may inhibit cell migration on Fibronectin.

REFERENCES

1. Jackson, T.R., et al. 2000. ACAPs are ARF6 GTPase-activating proteins that function in the cell periphery. *J. Cell Biol.* 151: 627-638.
2. Furman, C., et al. 2002. DEF-1/ASAP1 is a GTPase-activating protein (GAP) for ARF1 that enhances cell motility through a GAP-dependent mechanism. *J. Biol. Chem.* 277: 7962-7969.
3. Nie, Z., et al. 2003. Specific regulation of the adaptor protein complex AP-3 by the ARF GAP AGAP1. *Dev. Cell* 5: 513-521.
4. Dai, J., et al. 2004. ACAP1 promotes endocytic recycling by recognizing recycling sorting signals. *Dev. Cell* 7: 771-776.
5. Ivaska, J., et al. 2005. PKC ϵ -mediated phosphorylation of vimentin controls integrin recycling and motility. *EMBO J.* 24: 3834-3845.

CHROMOSOMAL LOCATION

Genetic locus: ACAP1 (human) mapping to 17p13.1.

SOURCE

ACAP1 (B-11) is a mouse monoclonal antibody raised against amino acids 18-172 mapping near the N-terminus of ACAP1 of human origin.

PRODUCT

Each vial contains 200 μ g IgG γ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

ACAP1 (B-11) is available conjugated to agarose (sc-376574 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-376574 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-376574 PE), fluorescein (sc-376574 FITC), Alexa Fluor[®] 488 (sc-376574 AF488), Alexa Fluor[®] 546 (sc-376574 AF546), Alexa Fluor[®] 594 (sc-376574 AF594) or Alexa Fluor[®] 647 (sc-376574 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-376574 AF680) or Alexa Fluor[®] 790 (sc-376574 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

ACAP1 (B-11) is recommended for detection of ACAP1 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for ACAP1 siRNA (h): sc-44442, ACAP1 shRNA Plasmid (h): sc-44442-SH and ACAP1 shRNA (h) Lentiviral Particles: sc-44442-V.


Molecular Weight of ACAP1: 80 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200 or Jurkat whole cell lysate: sc-2204.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA



ACAP1 (B-11): sc-376574. Western blot analysis of ACAP1 expression in Jurkat whole cell lysate.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.